



Apr 23rd, 9:00 AM - 4:00 PM

Exotic, Three Dimensional Macro-Structures in Gels Produced by Kinetically Limited and Geometrically Constrained Coarsening

Dana Deardorff

Illinois Wesleyan University

Kimberly Branshaw

Illinois Wesleyan University

Narendra K. Jaggi, Faculty Advisor

Illinois Wesleyan University

Follow this and additional works at: <http://digitalcommons.iwu.edu/jwprc>

Deardorff, Dana; Branshaw, Kimberly; and Jaggi, Faculty Advisor, Narendra K., "Exotic, Three Dimensional Macro-Structures in Gels Produced by Kinetically Limited and Geometrically Constrained Coarsening" (1994). *John Wesley Powell Student Research Conference*. 24.

<http://digitalcommons.iwu.edu/jwprc/1994/posters/24>

This Event is brought to you for free and open access by The Ames Library, the Andrew W. Mellon Center for Curricular and Faculty Development, the Office of the Provost and the Office of the President. It has been accepted for inclusion in Digital Commons @ IWU by the faculty at Illinois Wesleyan University. For more information, please contact digitalcommons@iwu.edu.

©Copyright is owned by the author of this document.

EXOTIC, THREE DIMENSIONAL MACRO-STRUCTURES IN
GELS PRODUCED BY KINETICALLY LIMITED AND
GEOMETRICALLY CONSTRAINED COARSENING

Dana Deardorff, Kimberly Branshaw and Narendra K. Jaggi *
Physics Department, Illinois Wesleyan University

We have recently discovered a rare and elegant coarsening mode in some polyelectrolyte gels that leads to intermediate structures that are complex and beautiful. They were discovered accidentally when a relatively large and thick cylindrical piece of the high density, crosslinked polymer was allowed to swell in water over an extended period. The resulting structure and its time evolution are not easily described in a capsule form. They evolve from monotonous cylindrical shapes in the beginning to periodic, circular, gear-tooth-like structures at short times. At intermediate stages, the elegant three dimensional, orchid-like patterns border on the sensuous. After a long time (of the order of days for length scales of an inch or so), the object is a scaled-up replica of the initial cylindrical structure. Curiously, the intermediate structures, however complex, are quite reproducible in all essential features. The results and our qualitative understanding of this phenomenon will be presented.

Acknowledgment: This research was supported in part by a grant (# NAG-8-258) from NASA under the NASA/JOVE program