

## Illinois Wesleyan University Digital Commons @ IWU

John Wesley Powell Student Research Conference

1994, 5th Annual JWP Conference

Apr 23rd, 9:00 AM - 4:00 PM

## The Recurrence Relation of B-Wavelets

Rumi Kumazawa, '94 Illinois Wesleyan University

Tian-Xiao He, Faculty Advisor Illinois Wesleyan University

Follow this and additional works at: http://digitalcommons.iwu.edu/jwprc

Rumi Kumazawa, '94 and Tian-Xiao He, Faculty Advisor, "The Recurrence Relation of B-Wavelets" (April 23, 1994). *John Wesley Powell Student Research Conference*. Paper 29. http://digitalcommons.iwu.edu/jwprc/1994/posters/29

This Event is brought to you for free and open access by the Conferences & Events at Digital Commons @ IWU. It has been accepted for inclusion in John Wesley Powell Student Research Conference by an authorized administrator of Digital Commons @ IWU. For more information, please contact sdaviska@iwu.edu.

©Copyright is owned by the author of this document.

## THE RECURRENCE RELATION OF B-WAVELETS

Rumi Kumazawa and Tian-Xiao He\*, Department of Mathematics, IWU

The study of wavelet functions is a relatively new area in mathematics. It is a topic of interest for both mathematicians and engineers, and applications can be seen in a wide area where Fourier transforms were used traditionally.

The wavelet functions can be used to perform the decomposition in  $L^2$  space, the collection of all square integrable functions. This can be done with the aid of compact support functions called the B-spline functions. The B-splines act as scaling functions in order to construct specific wavelet functions - the B-wavelets.

B-spline functions as piecewise polynomials with compact supports can be relatively smooth, and their Fourier transforms possess some properties like the Dirac Delta functions. Thus B-wavelets as the dilations and translations of B-splines can be used to reproduce and to analyze signals both in the local time and local frequency domains. In addition, we can expect to find a recurrence relation of the B-spline functions with different order. Hence B-wavelets of any order can be constructed successively from the lower order ones.

My project is on constructing B-spline functions and B-wavelet functions by their recurrence relations. A program was written in Pascal in order to calculate the Bernstein-Bezier coefficients of the B-spline functions of different order, and the graphs of these functions were drawn using Mathematica. From these coefficients of the B-splines, the coefficients of corresponding B-wavelength functions were found.