



Apr 23rd, 9:00 AM - 4:00 PM

Toward an Understanding of Alzheimer's Disease I: The Effects of Bilateral Injections of Beta Amyloid (1-42) on Spatial Learning in the Male Rat

Anthony R. Peterson
Illinois Wesleyan University

Wayne Dornan, Faculty Advisor
Illinois Wesleyan University

Follow this and additional works at: <https://digitalcommons.iwu.edu/jwprc>

Peterson, Anthony R. and Dornan, Faculty Advisor, Wayne, "Toward an Understanding of Alzheimer's Disease I: The Effects of Bilateral Injections of Beta Amyloid (1-42) on Spatial Learning in the Male Rat" (1994). *John Wesley Powell Student Research Conference*. 42. <https://digitalcommons.iwu.edu/jwprc/1994/posters/42>

This is protected by copyright and/or related rights. It has been brought to you by Digital Commons @ IWU with permission from the rights-holder(s). You are free to use this material in any way that is permitted by the copyright and related rights legislation that applies to your use. For other uses you need to obtain permission from the rights-holder(s) directly, unless additional rights are indicated by a Creative Commons license in the record and/ or on the work itself. This material has been accepted for inclusion by faculty at Illinois Wesleyan University. For more information, please contact digitalcommons@iwu.edu.

©Copyright is owned by the author of this document.

TOWARD AN UNDERSTANDING OF ALZHEIMER'S DISEASE I:
THE EFFECTS OF BILATERAL INJECTIONS OF BETA AMYLOID (1-42)
ON SPATIAL LEARNING IN THE MALE RAT

Anthony R. Peterson and Wayne Doman*, Department of Psychology, IWU

Alzheimer's Disease is a neurodegenerative disorder that is characterized by neurofibrillary tangles (NFT) and neuritic plaques (NP). These NP are comprised of a 39 - 43 amino acid peptide : beta amyloid (β A), which seem to selectively target the basal forebrain and hippocampus of the brain. A recent study by Kang, Doman, McCampbell, Kang (Neuroreport, 1993) stated that β A(25-35), the hypothesized neurotoxic segment of the protein, plus a subthreshold dose of an excitatory amino acid, ibotenic acid (IBO), produced behavioral deficits on a visual-spatial task in a rat similar to the behavioral deficits in human Alzheimer's subjects. The same study reported that β A(25-35) alone did not produce any behavioral deficits. Presently there are several forms of β A that are suspected of forming NP. Research on the effects of these different fragments of β A is needed to discover their potential role in the etiology of Alzheimer's. Therefore, in this study, we assessed the behavioral effects of β A(1-42) following intracerebral injections into the dorsal blade (CA1 region) of the hippocampus in the male rat. We used three groups of animals in this study. One group consisted of animals injected with β A(1-42) in a dimethylsulfoxide vehicle (DMSO). The second group was comprised of animals that were injected with a scrambled β A(1-42) peptide in DMSO. The third group, the control group, consisted of animals injected with DMSO only. All of the injections were targeted at the dorsal blade of the hippocampus. The animals were pretested for two weeks on an eight arm radial arm maze, a task that tests for visual-spatial learning, in which five of the arms were always baited. We tested the animals for two weeks after the surgery and found no difference between any of the groups in their ability to learn a visual-spatial task. Another study is currently examining the effects of β A(1-42) with IBO.