Illinois Wesleyan University
Digital Commons @ IWU

John Wesley Powell Student Research Conference

2004, 15th Annual JWP Conference

Apr 17th, 1:15 PM - 2:30 PM

Analysis of the Large Amplitude Pendulum and the Effects of Damping

Jason Callard
Illinois Wesleyan University

Jason Dulac
Illinois Wesleyan University

William Brandon, Faculty Advisor
Illinois Wesleyan University

Follow this and additional works at: http://digitalcommons.iwu.edu/jwprc

Jason Callard; Jason Dulac; and William Brandon, Faculty Advisor, "Analysis of the Large Amplitude Pendulum and the Effects of Damping" (April 17, 2004). *John Wesley Powell Student Research Conference.* Paper 7.
http://digitalcommons.iwu.edu/jwprc/2004/posters2/7

This Event is brought to you for free and open access by The Ames Library, the Andrew W. Mellon Center for Curricular and Faculty Development, the Office of the Provost and the Office of the President. It has been accepted for inclusion in Digital Commons @ IWU by the faculty at Illinois Wesleyan University. For more information, please contact digitalcommons@iwu.edu.
©Copyright is owned by the author of this document.

Poster Presentation P14

ANALYSIS OF THE LARGE AMPLITUDE PENDULUM AND THE EFFECTS OF DAMPING

Jason Callard, Jason Dulac and William Brandon*
Department of Physics, Illinois Wesleyan University

We set out to study how a real pendulum oscillates about its equilibrium. Unlike the idealized pendulum, the period of oscillation is dependent upon the angle from which the pendulum is released and damps (decreases) with each oscillation. We found both period and damping to be dependent upon the angle from which the pendulum is released. In our case, the condition is accurately represented by Bernoulli's equation to the fourth order $T = 2\pi\sqrt{\frac{L}{g}}\left(1+\frac{1}{4}\sin^2\frac{\theta}{2}+\frac{9}{64}\sin^4\frac{\theta}{2}\right)$, where L is the length to the center of mass and θ is the angle of release as measured from the perpendicular. Due to the variance in non-conservative forces, there is no explicit equation to apply to damping because all damping equations rely on measured constants. However, the extent of damping increased as θ increased, and we were able to depict damping for our particular pendulum.