



Apr 16th, 1:15 PM - 2:30 PM

Stabilizing A [4FE-4S] Cluster to Oxygen: Effects of Amino Acid Substitutions Near the Cysteine Ligands in the FNR Transcription Factor

Robin D. Johnson

Illinois Wesleyan University

Lindsay A. Sicks

Illinois Wesleyan University

Maura K. Bates

Illinois Wesleyan University

Erin N. Fleck, '06

Illinois Wesleyan University

Laura Moore, Faculty Advisor

Illinois Wesleyan University

Follow this and additional works at: <http://digitalcommons.iwu.edu/jwprc>

Johnson, Robin D.; Sicks, Lindsay A.; Bates, Maura K.; Fleck, '06, Erin N.; and Moore, Faculty Advisor, Laura, "Stabilizing A [4FE-4S] Cluster to Oxygen: Effects of Amino Acid Substitutions Near the Cysteine Ligands in the FNR Transcription Factor" (2005). *John Wesley Powell Student Research Conference*. 3.
<http://digitalcommons.iwu.edu/jwprc/2005/posters2/3>

This Event is brought to you for free and open access by The Ames Library, the Andrew W. Mellon Center for Curricular and Faculty Development, the Office of the Provost and the Office of the President. It has been accepted for inclusion in Digital Commons @ IWU by the faculty at Illinois Wesleyan University. For more information, please contact digitalcommons@iwu.edu.

©Copyright is owned by the author of this document.

Poster Presentation P8

STABILIZING A [4Fe-4S] CLUSTER TO OXYGEN: EFFECTS OF AMINO ACID SUBSTITUTIONS NEAR THE CYSTEINE LIGANDS IN THE FNR TRANSCRIPTION FACTOR

Robin D. Johnson, Lindsay A. Sicks, Maura K. Bates, and Erin N. Fleck
and Laura Moore*

Chemistry/Biology Departments, Illinois Wesleyan University

The FNR protein is an oxygen-sensing transcription factor found in the facultative anaerobe *Escherichia coli*. Dimerization and DNA binding of wild type FNR are coupled to the incorporation of a [4Fe-4S] cluster, which is ligated by cysteine residues at positions 20, 23, 29, and 122.

In the presence of molecular oxygen, this cluster degrades causing a conformational change that renders the protein inactive. Previous research has shown that the Leu28His mutant FNR protein has an oxygen stable [4Fe-4S] cluster. Several more amino acid substitutions have been made at residues adjacent to the cysteine ligands. The in vivo activity of these mutant FNR proteins has been evaluated by beta-galactosidase assays under aerobic and anaerobic conditions. Most of the mutant proteins retained similar activity to that of the wild type protein, with significant activity under anaerobic condition and little activity under aerobic conditions. However, replacement of the Leu28 with Lys or Arg resulted in proteins that are active under aerobic conditions.

These results suggest that a basic amino acid residue at position 28 stabilizes the cluster to oxygen. The Leu28Lys FNR protein has been isolated and preliminary characterization of its [4Fe-4S] cluster by absorption spectroscopy shows that like the Leu28His FNR protein, the [4Fe-4S] cluster in Leu28Lys FNR is stable in the presence of oxygen.

The [4Fe-4S] cluster of the Leu28Lys mutant protein is currently being further characterized in order to give a better understanding of how a basic amino acid at position 28 helps to stabilize the cluster in the presence of oxygen.