



Apr 17th, 1:15 PM - 2:30 PM

The Effect of Light on Rates of Cloning of the Symbiont-Bearing Acoel *Convolutriloba Longifissura*

Sara Rahim

Illinois Wesleyan University

William Jaeckle, Faculty Advisor

Illinois Wesleyan University

Follow this and additional works at: <http://digitalcommons.iwu.edu/jwprc>

Rahim, Sara and Jaeckle, Faculty Advisor, William, "The Effect of Light on Rates of Cloning of the Symbiont-Bearing Acoel *Convolutriloba Longifissura*" (2004). *John Wesley Powell Student Research Conference*. 21.
<http://digitalcommons.iwu.edu/jwprc/2004/posters2/21>

This Event is brought to you for free and open access by The Ames Library, the Andrew W. Mellon Center for Curricular and Faculty Development, the Office of the Provost and the Office of the President. It has been accepted for inclusion in Digital Commons @ IWU by the faculty at Illinois Wesleyan University. For more information, please contact digitalcommons@iwu.edu.

©Copyright is owned by the author of this document.

Poster Presentation P42

**THE EFFECT OF LIGHT ON RATES OF CLONING OF THE
SYMBIONT-BEARING ACOEL *CONVOLUTRILOBA LONGIFISSURA***

Sara Rahim and William Jaeckle*
Department of Biology, Illinois Wesleyan University

The acoel turbellarian *Convolutriloba longifissura* reproduces primarily by asexual fission and engages in an obligate symbiotic relationship with unicellular algae belonging to the genus *Tetraselmis*. The obligate nature of the symbiosis between these species suggests that algal photosynthesis may influence rates of flatworm asexual reproduction. To test this hypothesis we explored the effect of light on *C. longifissura's* ability to clone. Worms (n=18/treatment) were incubated at light regimes of total darkness, 8L:16D, and 12L:12D for 30 days at $26 \pm 1^\circ\text{C}$. Worms held in complete darkness experienced 100% mortality after 14 days; mortality in all other light treatments was <4%. Cloning increased with a greater exposure to light (p=0.03). The average rate of cloning for worms exposed to 8L:16D light cycle was 0.022/day; rates of cloning by individuals held at 12D:12L was nearly 3X greater (0.061/day). Flatworm length and cloning rate were positively correlated ($r=0.77$, $p=0.01$, $n=36$); there were no significant differences in worm length among light treatments ($p>0.05$). Differences were detected in the cloning rates of worms exposed to blue, red, green, or white light for 20 days at $26 \pm 1^\circ\text{C}$. Cloning rates were highest for worms exposed to white light, lowest for worms exposed to green light, and intermediate for those exposed to red or blue light. These data provide evidence supporting the hypothesis that algal photosynthetic activity directly affects cloning by *C. longifissura*. We suggest that light conditions which promote photosynthesis in algae result in release of photosynthate to the flatworm and these materials fuel asexual reproduction. In turn, flatworms provide shelter and/or nutrients to the algae. This allows for the possibility of a mutualistic symbiotic relationship between *C. longifissura* and *Tetraselmis* sp.