

Illinois Wesleyan University Digital Commons @ IWU

John Wesley Powell Student Research Conference

2003, 14th Annual JWP Conference

Apr 12th, 1:15 PM - 2:15 PM

Organochlorine Pesticide Contamination Patterns in Nearctic Resident Birds

Jeffrey Horvath, '03 Illinois Wesleyan University

R. Given Harper, Faculty Advisor Illinois Wesleyan University

Follow this and additional works at: http://digitalcommons.iwu.edu/jwprc

Jeffrey Horvath, '03 and R. Given Harper, Faculty Advisor, "Organochlorine Pesticide Contamination Patterns in Nearctic Resident Birds" (April 12, 2003). *John Wesley Powell Student Research Conference*. Paper 16. http://digitalcommons.iwu.edu/jwprc/2003/posters2/16

This Event is brought to you for free and open access by the Conferences & Events at Digital Commons @ IWU. It has been accepted for inclusion in John Wesley Powell Student Research Conference by an authorized administrator of Digital Commons @ IWU. For more information, please contact sdaviska@iwu.edu.

©Copyright is owned by the author of this document.

Poster Presentation P18

ORGANOCHLORINE PESTICIDE CONTAMINATION PATTERNS IN NEARCTIC RESIDENT BIRDS

<u>Jeffrey Horvath</u> and R. Given Harper* Department of Biology, Illinois Wesleyan University

Thirteen species of Nearctic resident passerines and woodpeckers collected in central Illinois and North Carolina between 1991 and 2001 were analyzed for the presence of 17 organochlorine (OC) pesticides and metabolites. OC compound residues were detected in 29 of the 32 individuals examined.

Contaminated birds contained from one to eleven different compounds, with aldrin (mean \pm SE; 97.25 \pm 35.16 ng/g), beta-BHC (20.60 \pm 5.36 ng/g), DDE (26.65 \pm 4.53 ng/g), dieldrin (24.88 \pm 7.99 ng/g), endosulfan I (24.13 \pm 4.76 ng/g), heptachlor (13.02 \pm 2.40 ng/g), and heptachlor epoxide (18.35 \pm 2.27 ng/g) appearing most frequently. Levels of contamination for specific compounds ranged from 3.84 ng/g to 688.95 ng/g. For the 15 specimens from Illinois, there was no significant difference in total OC levels in birds collected in urban locations (370.50 \pm 143.43 ng/g) compared to birds collected in rural locations (112.91 \pm 25.38 ng/g; t = -1.95, df = 10.63, P = 0.08). There was no significant difference in total OCs in birds collected from Illinois vs. North Carolina. Likewise, in the pooled sample there was no significant effect of diet (granivorous/insectivorous vs. insectivorous), or age class (AHY vs. HY) on total pesticide levels.

Use and production of the majority of the OC compounds detected have been banned or restricted in North America for 5-30 years. The high frequency of OC contamination in Nearctic resident birds indicates the presence of OC compounds in the environment through extended persistence, atmospheric deposition or illegal usage.

Jeffrey J. Horvath¹, R. Given Harper¹, Jeffrey A. Frick², Angelo P. Capparella³, John Gerwin⁴, and Rebecca Browning⁴

¹Department of Biology, Illinois Wesleyan University, Bloomington, Illinois, USA 61702-2900

²Department of Chemistry, Illinois Wesleyan University, Bloomington, Illinois, USA 61702-2900

³Ecology Group, Department of Biological Sciences, Illinois State University, Normal, Illinois, USA 61790-4120

⁴North Carolina Museum of Natural Sciences, Raleigh, North Carolina, USA 27601-1029