



Apr 14th, 2:35 PM - 3:35 PM

# Creation of Stable Porphobilinogen Synthase Knockouts in E. Coli

Steven Cherney, '07

*Illinois Wesleyan University*

David Bollivar, Faculty Advisor

*Illinois Wesleyan University*

Follow this and additional works at: <http://digitalcommons.iwu.edu/jwprc>

Cherney, '07, Steven and Bollivar, Faculty Advisor, David, "Creation of Stable Porphobilinogen Synthase Knockouts in E. Coli" (2007). *John Wesley Powell Student Research Conference*. 10.  
<http://digitalcommons.iwu.edu/jwprc/2007/posters2/10>

This Event is brought to you for free and open access by The Ames Library, the Andrew W. Mellon Center for Curricular and Faculty Development, the Office of the Provost and the Office of the President. It has been accepted for inclusion in Digital Commons @ IWU by the faculty at Illinois Wesleyan University. For more information, please contact [digitalcommons@iwu.edu](mailto:digitalcommons@iwu.edu).

©Copyright is owned by the author of this document.

Poster Presentation P20

**CREATION OF STABLE PORPHOBILINOGEN SYNTHASE KNOCKOUTS IN E. COLI**

Steven Cherney and David Bollivar\*  
Biology Department, Illinois Wesleyan University

Porphobilinogen synthase (PBGs) is the enzyme responsible for the first step in the biosynthesis of tetrapyrroles such as chlorophyll and porphyrin. Our goal is to create a stable porphobilinogen synthase (PBGs) knockout in *Escherichia coli*. *Escherichia coli* does not have a natural system to import heme, the end product of the biochemical pathway in this organism. Consequently we also must also have a stable heme uptake system. Using a strain we obtained from another laboratory that has a heme uptake protein from another bacterium *S. marcescens*, we used PCR products to knockout the PBGS gene, and isolate organisms that must uptake heme from their environment. The PBGS knockout strain is being created so that the genes for PBGS from other organisms can be introduced and studied in an in vivo system.