

Illinois Wesleyan University Digital Commons @ IWU

John Wesley Powell Student Research Conference

2007, 18th Annual JWP Conference

Apr 14th, 2:35 PM - 3:35 PM

Variation In Metabolic Cost of Embryonic Development of the Freshwater Snail, Phsya SP.

Andrew Hall, '07 Illinois Wesleyan University

William Jaeckle, Faculty Advisor Illinois Wesleyan University

Follow this and additional works at: http://digitalcommons.iwu.edu/jwprc

Andrew Hall, '07 and William Jaeckle, Faculty Advisor, "Variation In Metabolic Cost of Embryonic Development of the Freshwater Snail, Phsya SP." (April 14, 2007). *John Wesley Powell Student Research Conference*. Paper 39. http://digitalcommons.iwu.edu/jwprc/2007/posters2/39

This Event is brought to you for free and open access by the Conferences & Events at Digital Commons @ IWU. It has been accepted for inclusion in John Wesley Powell Student Research Conference by an authorized administrator of Digital Commons @ IWU. For more information, please contact sdaviska@iwu.edu.

©Copyright is owned by the author of this document.

Poster Presentation P38

VARIATION IN METABOLIC COST OF EMBRYONIC DEVELOPMENT OF THE FRESHWATER SNAIL, PHYSA SP.

<u>Andrew Hall</u> and William Jaeckle* Biology Department, Illinois Wesleyan University

Embryos of the freshwater snail Physa sp. complete their development within an egg capsule and hatch as juveniles. To estimate the energetic cost of development, oxygen consumption rates of egg masses were monitored from deposition to hatching. Oxygen consumption increased during development (r2=0.173, p<0.01), but there was no consistent pattern. A negative correlation existed (r=-0.984, p<0.01) between hatching time and oxygen consumption, and a positive correlation existed (r=0.903, p<0.05) between the rate of oxygen consumption and total energy expended. Masses with longer times to hatching utilize less energy than masses with shorter times to hatching. If egg capsules are initially identical, then embryos with higher metabolic rates and shorter development times will be smaller at hatching than individuals with lower rates of energy consumption. If capsule size varies, then differences in metabolism may result in different hatching times, but a common juvenile size. The effects of these differences on juvenile survivorship in the field are unknown.