

Illinois Wesleyan University Digital Commons @ IWU

John Wesley Powell Student Research Conference

2012, 23rd Annual JWP Conference

Apr 14th, 10:00 AM - 11:00 AM

Comparison of Regional Eggshell Porosity Between the Brood Parasitic Brown-Headed Cowbird (*Moluthrus ater*) and its Hosts: the Dickcissel (*Spiza americana*), and Two Non-Parasitic Relatives, the Red-Winged Blackbird (*Agelaius phoeniceus*) and the Common Grackle (*Quiscalus quiscula*).

Brittany Childs *Illinois Wesleyan University*

W. B. Jaeckle, Faculty Advisor *Illinois Wesleyan University*

R. G. Harper, Faculty Advisor *Illinois Wesleyan University*

M. Kiefer *Illinois Wesleyan University*

Brittany Childs; W. B. Jaeckle, Faculty Advisor; R. G. Harper, Faculty Advisor; M. Kiefer; J. W. Rivers; and B. D. Peer, "Comparison of Regional Eggshell Porosity Between the Brood Parasitic Brown-Headed Cowbird (*Moluthrus ater*) and its Hosts: the Dickcissel (*Spiza americana*), and Two Non-Parasitic Relatives, the Red-Winged Blackbird (*Agelaius phoeniceus*) and the Common Grackle (*Quiscalus quiscula*)." (April 14, 2012). *John Wesley Powell Student Research Conference*. Paper 3.

http://digitalcommons.iwu.edu/jwprc/2012/oralpres/3

This Event is brought to you for free and open access by The Ames Library, the Andrew W. Mellon Center for Curricular and Faculty Development, the Office of the Provost and the Office of the President. It has been accepted for inclusion in Digital Commons @ IWU by the faculty at Illinois Wesleyan University. For more information, please contact digitalcommons@iwu.edu.

©Copyright is owned by the author of this document.

J. W. Rivers

Oregon State University

See next page for additional authors

Follow this and additional works at: http://digitalcommons.iwu.edu/jwprc Part of the <u>Biology Commons</u>



Oral Presentation O1.3

COMPARISON OF REGIONAL EGGSHELL POROSITY BETWEEN THE BROOD PARASITIC BROWN-HEADED COWBIRD (MOLUTHRUS ATER) AND ITS HOSTS: THE DICKCISSEL (SPIZA AMERICANA), AND TWO NON-PARASITIC RELATIVES, THE RED-WINGED BLACKBIRD (AGELAIUS PHOENICEUS) AND THE COMMON GRACKLE (QUISCALUS QUISCULA).

Brittany Childs^a, W.B. Jaeckle^{a*}, R.G. Harper^{a*}, M. Kiefer^a, J.W. Rivers^b, B.D. Peer^c

^aDepartment of Biology, Illinois Wesleyan University, Bloomington, IL

^bDepartment of Forest Ecosystems and Society, Oregon State University, Corvallis, OR

^cDepartment of Biological Sciences, Western Illinois University, Macomb, IL

The Brown-headed Cowbird is a generalist brood parasite that lays eggs in the nests of many host species, including the Dickcissel and two non-parasitic relatives: the Red-winged Blackbird and the Common Grackle. Cowbird eggs reportedly hatch sooner than equivalently-sized host eggs, presumably via accelerated embryonic development enabled by a greater eggshell porosity and consequently greater gas exchange. The distribution of pores among apical, equatorial and basal eggshell regions within cowbirds and host species is undetermined. We tested the hypothesis that equatorial porosity would be greatest because respiratory gases cross the eggshell and enter or exit the embryo's circulatory system in that region. We found that, when normalized to mass, cowbird eggshells had significantly greater pore area and porosity in equatorial regions compared to its three hosts ($p \le 0.001$). Cowbird eggshells had greater apical pore area than the hosts (p < 0.04) but equivalent basal pore area (p > 0.09) and porosity (p > 0.15) to its relatives.