Illinois Wesleyan University
Digital Commons @ IWU

John Wesley Powell Student Research Conference

2012, 23rd Annual JWP Conference

Apr 14th, 2:35 PM - 3:35 PM

Metabolic Contribution of Particulate Matter (<μm) for Larvae of *Lytechinus variegatus*

Matthew McShane '13
Illinois Wesleyan University

William Jaeckle, Faculty Advisor
Illinois Wesleyan University

Follow this and additional works at: http://digitalcommons.iwu.edu/jwprc

Matthew McShane '13 and William Jaeckle, Faculty Advisor, "Metabolic Contribution of Particulate Matter (<μm) for Larvae of *Lytechinus variegatus*" (April 14, 2012). *John Wesley Powell Student Research Conference.* Paper 12.
http://digitalcommons.iwu.edu/jwprc/2012/posters2/12

This Event is brought to you for free and open access by The Ames Library, the Andrew W. Mellon Center for Curricular and Faculty Development, the Office of the Provost and the Office of the President. It has been accepted for inclusion in Digital Commons @ IWU by the faculty at Illinois Wesleyan University. For more information, please contact digitalcommons@iwu.edu.
©Copyright is owned by the author of this document.

Poster Presentation P24

METABOLIC CONTRIBUTION OF PARTICULATE MATTER (<1µm) FOR LARVAE OF *LYTECHINUS VARIEGATUS*

Matthew McShane and William Jaeckle*
Biology Department, Illinois Wesleyan University

Echinoderm larvae possess an external band of cilia that beat in one direction for locomotion and undergo localized beat reversal when triggered by particles (≥ 2 mm), pushing these particles towards the mouth for ingestion. Particles below adequate size, if ingested, must be collected by passing seawater through the digestive system. We measured the ability of pluteus larvae of *Lytechinus variegatus* to capture bacteria-sized particles after exposing them to 0.513µm fluorescent beads (9.5×10^5 beads/mL) for 10 min. Beads were found in the digestive system in 75% (41 of 55) of larvae and the average ingestion rate was 39.4 (±42.7 SD) beads/h. From these ingestion rates we calculated clearance rates (mL/ larva-h) and used literature values for bacterial energy content and larval metabolism to estimate the energetic contribution of bacteria-sized particles. Assuming 100% assimilation efficiency and 10^6 cells/mL, we found that feeding on bacteria-sized particles could provide for only 0.16% of metabolic rate.