



Illinois Wesleyan University
Digital Commons @ IWU

John Wesley Powell Student Research
Conference

1990, 1st Annual JWP Conference

Apr 6th, 6:30 PM - 8:00 PM

Bilateral Ibotenic Acid Lesions of the Bednucleus of the Stria Terminalis in Intact Male Rats: a Pilot Study

Christopher Ballak
Illinois Wesleyan University

Dr. Wayne Dornan, Faculty Advisor
Illinois Wesleyan University

Follow this and additional works at: <https://digitalcommons.iwu.edu/jwprc>

Ballak, Christopher and Dornan, Faculty Advisor, Dr. Wayne, "Bilateral Ibotenic Acid Lesions of the Bednucleus of the Stria Terminalis in Intact Male Rats: a Pilot Study" (1990). *John Wesley Powell Student Research Conference*. 1.
<https://digitalcommons.iwu.edu/jwprc/1990/posters/1>

This Event is protected by copyright and/or related rights. It has been brought to you by Digital Commons @ IWU with permission from the rights-holder(s). You are free to use this material in any way that is permitted by the copyright and related rights legislation that applies to your use. For other uses you need to obtain permission from the rights-holder(s) directly, unless additional rights are indicated by a Creative Commons license in the record and/ or on the work itself. This material has been accepted for inclusion by faculty at Illinois Wesleyan University. For more information, please contact digitalcommons@iwu.edu.

©Copyright is owned by the author of this document.

BILATERAL IBOTENIC ACID LESIONS OF THE BED NUCLEUS OF THE STRIA TERMINALIS IN INTACT MALE RATS: A PILOT STUDY

Christopher Ballak, Dept. of Psychology, IWU, Dr. Wayne Dornan*

The bed nucleus of the stria terminalis (BnST) is commonly thought of as a "relay station" integrating vomeronasal and olfactory information from the medial nucleus of the amygdala (MeA) to the medial preoptic area (MPOA). Recently, Malsbury and Mc Kay (Brain Research Bulletin, '89) reported a sex difference in the pattern of the substance P-immunoreactive (sP-ir) innervation of the medial division of the BnST (larger in males). Moreover, they suggest that the sexually dimorphic innervation of the medial BnST originates from the MeA. Interestingly, sP has been reported to facilitate male sexual behavior when bilaterally injected into the MPOA. Taken together, this suggests that the sP innervation of the BnST may be important in the regulation of male sexual behavior and that the BnST might be more than a relay station for the MeA. Numerous sP-ir cell bodies are present in the BnST, presently, however, nothing is known concerning whether this sP cell group is necessary for the expression of male sexual behavior. Therefore, the purpose of this study is two fold: First, we will examine the effects of ibotenic acid lesions of the BnST on male sexual behavior in order to determine if the BnST is only a relay station or if it is involved in the regulation of male sexual behavior. Second, following the destruction of cell bodies of the medial BnST, we will have effectively eliminated the influence of the sP sexually dimorphic innervation of the medial BnST this will elucidate whether this sP innervation is involved in the neural regulation of sexual behavior in the male rat. The results of this study will further the understanding of the role of sP in sexual behavior.

Adult Long-Evans male rats will be used. Each male will receive either a bilateral 0.3ul pressure injection of either ibotenic acid (10ug/1ul) or phosphate buffered saline into the medial BnST (AP = -0.82; ML = -1.2; DV = -7.1). All males will be tested prior to surgery and then again five - seven days following surgery. A variety of parameters of male sexual behavior will be measured. We postulate that an inhibition of male sexual behavior will be observed.