



Illinois Wesleyan University
Digital Commons @ IWU

John Wesley Powell Student Research
Conference

1992, 3rd Annual JWP Conference

Apr 25th, 10:30 AM - 4:30 PM

Analysis of Carbon across the Cenomanian-Turonian Boundary

Andrew J. Carr
Illinois Wesleyan University

Wendy S. Wolbach, Faculty Advisor
Illinois Wesleyan University

Follow this and additional works at: <https://digitalcommons.iwu.edu/jwprc>

Carr, Andrew J. and Wolbach, Faculty Advisor, Wendy S., "Analysis of Carbon across the Cenomanian-Turonian Boundary" (1992). *John Wesley Powell Student Research Conference*. 45.

<https://digitalcommons.iwu.edu/jwprc/1992/posters/45>

This Event is protected by copyright and/or related rights. It has been brought to you by Digital Commons @ IWU with permission from the rights-holder(s). You are free to use this material in any way that is permitted by the copyright and related rights legislation that applies to your use. For other uses you need to obtain permission from the rights-holder(s) directly, unless additional rights are indicated by a Creative Commons license in the record and/ or on the work itself. This material has been accepted for inclusion by faculty at Illinois Wesleyan University. For more information, please contact digitalcommons@iwu.edu.

©Copyright is owned by the author of this document.

ANALYSIS OF CARBON ACROSS THE CENOMANIAN-TURONIAN BOUNDARY

Andrew J. Carr and Wendy S. Wolbach*
Dept of Chemistry, Illinois Wesleyan University

Soot from wildfires has been discovered in sedimentary rocks at the 66 Ma-old Cretaceous-Tertiary (K-T) boundary, a time of mass extinctions which included those of the dinosaurs. This extinction was presumably caused by the impact of a giant meteorite. The question at hand is whether other extinctions have been caused by meteorite impacts as well. Discovery of impact-related evidence at a smaller extinction event that occurred during the Cenomanian-Turonian boundary (C-T) 92 Ma ago, where plankton were the chief casualty, could lend support to the impact theory. Analysis of sedimentary rocks spanning this extinction horizon for possible changes in abundance, morphology, and isotopic composition of reduced organic and inorganic carbon (soot) may yield such information. In the absence of soot, the isotopic study of the organic carbon could yield important clues regarding changes in the environment at that time.

Currently two sample sites are being analyzed for reduced carbon content: Red Wash, New Mexico and Chipsa Summit, Texas. Both sample sites are from the Western Interior Basin. During the time of the extinctions 92 million years ago, the basin was an inland sea. The preliminary data suggests a decrease in the amount of reduced carbon at both sample sites. This would indicate that the extinction that took place was not rapid. Lower amounts of reduced carbon are generally more characteristic of extinctions caused by changes in climate conditions, compared with those caused by meteorite impacts. Climate changes would allow microorganisms time to digest (oxidize) the plankton before sedimentation could occur, thus reducing the amount of reduced carbon preserved in the rock.