



Apr 25th, 10:30 AM - 4:30 PM

Analysis of Carbon across the Cenomanian-Turonian Boundary

Andrew J. Carr

Illinois Wesleyan University

Wendy S. Wolbach, Faculty Advisor

Illinois Wesleyan University

Follow this and additional works at: <http://digitalcommons.iwu.edu/jwprc>

Carr, Andrew J. and Wolbach, Faculty Advisor, Wendy S., "Analysis of Carbon across the Cenomanian-Turonian Boundary" (1992). *John Wesley Powell Student Research Conference*. 45.
<http://digitalcommons.iwu.edu/jwprc/1992/posters/45>

This Event is brought to you for free and open access by The Ames Library, the Andrew W. Mellon Center for Curricular and Faculty Development, the Office of the Provost and the Office of the President. It has been accepted for inclusion in Digital Commons @ IWU by the faculty at Illinois Wesleyan University. For more information, please contact digitalcommons@iwu.edu.

©Copyright is owned by the author of this document.

ANALYSIS OF CARBON ACROSS THE CENOMANIAN-TURONIAN BOUNDARY

Andrew J. Carr and Wendy S. Wolbach*
Dept of Chemistry, Illinois Wesleyan University

Soot from wildfires has been discovered in sedimentary rocks at the 66 Ma-old Cretaceous-Tertiary (K-T) boundary, a time of mass extinctions which included those of the dinosaurs. This extinction was presumably caused by the impact of a giant meteorite. The question at hand is whether other extinctions have been caused by meteorite impacts as well. Discovery of impact-related evidence at a smaller extinction event that occurred during the Cenomanian-Turonian boundary (C-T) 92 Ma ago, where plankton were the chief casualty, could lend support to the impact theory. Analysis of sedimentary rocks spanning this extinction horizon for possible changes in abundance, morphology, and isotopic composition of reduced organic and inorganic carbon (soot) may yield such information. In the absence of soot, the isotopic study of the organic carbon could yield important clues regarding changes in the environment at that time.

Currently two sample sites are being analyzed for reduced carbon content: Red Wash, New Mexico and Chipsa Summit, Texas. Both sample sites are from the Western Interior Basin. During the time of the extinctions 92 million years ago, the basin was an inland sea. The preliminary data suggests a decrease in the amount of reduced carbon at both sample sites. This would indicate that the extinction that took place was not rapid. Lower amounts of reduced carbon are generally more characteristic of extinctions caused by changes in climate conditions, compared with those caused by meteorite impacts. Climate changes would allow microorganisms time to digest (oxidize) the plankton before sedimentation could occur, thus reducing the amount of reduced carbon preserved in the rock.