



Illinois Wesleyan University
Digital Commons @ IWU

John Wesley Powell Student Research
Conference

1993, 4th Annual JWP Conference

May 8th, 9:30 AM - 4:30 PM

Effect of Temperature on Growth of the Zebra Mussel, *Dreissena Polymorpha*

Tamara K. Ross
Illinois Wesleyan University

Gail Lima, Faculty Advisor
Illinois Wesleyan University

Follow this and additional works at: <https://digitalcommons.iwu.edu/jwprc>

Ross, Tamara K. and Lima, Faculty Advisor, Gail, "Effect of Temperature on Growth of the Zebra Mussel, *Dreissena Polymorpha*" (1993). *John Wesley Powell Student Research Conference*. 12.

<https://digitalcommons.iwu.edu/jwprc/1993/posters/12>

This Event is protected by copyright and/or related rights. It has been brought to you by Digital Commons @ IWU with permission from the rights-holder(s). You are free to use this material in any way that is permitted by the copyright and related rights legislation that applies to your use. For other uses you need to obtain permission from the rights-holder(s) directly, unless additional rights are indicated by a Creative Commons license in the record and/ or on the work itself. This material has been accepted for inclusion by faculty at Illinois Wesleyan University. For more information, please contact digitalcommons@iwu.edu.

©Copyright is owned by the author of this document.

EFFECT OF TEMPERATURE ON GROWTH OF THE ZEBRA MUSSEL, *DREISSENA POLYMORPHA*

Tamara K. Ross, Dept. of Biology, IWU, Gail Lima*

Dreissena polymorpha (Pallas), zebra mussels, are recent invaders of North American freshwater systems. They have a high reproductive rate and settle in high densities which can clog water intake valves and pipes. Many studies investigating the use of heat as a control measure have examined the effects of high temperatures on zebra mussel mortality. Much less is known about the effect of temperature on the actual growth rate and development of zebra mussels. This study examined the growth rates of zebra mussels at 10°C, 20°C, and 25°C over two four-week periods in the laboratory. Mussels were placed in culture dishes (five similarly sized mussels per dish) and fed 100 ml of the algae *Chlorella pyrenoidosa* (Chick) daily at a concentration of 4.13×10^5 cells/ml. Shell length and shell height measurements were taken three times during the experiment. Although shell length is the measurement typically used in growth studies, it has not been documented whether shell length is a better growth indicator than shell height. Both shell length and shell height are currently being compared with tissue weight to see which is a more accurate measure of growth. The results of this analysis will then be used to determine what effect temperature has on the growth of the mussels. The findings of this study will be important because factors which reduce growth influence the longterm success of the zebra mussels and therefore should be useful as possible control methods. Also, scientists working with zebra mussels in the laboratory will find the results of this study useful in determining the best conditions in which to raise zebra mussels.