



Illinois Wesleyan University
Digital Commons @ IWU

John Wesley Powell Student Research
Conference

1994, 5th Annual JWP Conference

Apr 23rd, 9:00 AM - 4:00 PM

β -Amyloid (1-42) and Its Role in Developing an Animal Model of Alzheimer's Disease

Greg Tinkler
Illinois Wesleyan University

Dr. Wayne Dornan, Faculty Advisor
Illinois Wesleyan University

Follow this and additional works at: <https://digitalcommons.iwu.edu/jwprc>

Tinkler, Greg and Dornan, Faculty Advisor, Dr. Wayne, " β -Amyloid (1-42) and Its Role in Developing an Animal Model of Alzheimer's Disease" (1994). *John Wesley Powell Student Research Conference*. 56.

<https://digitalcommons.iwu.edu/jwprc/1994/posters/56>

This Event is protected by copyright and/or related rights. It has been brought to you by Digital Commons @ IWU with permission from the rights-holder(s). You are free to use this material in any way that is permitted by the copyright and related rights legislation that applies to your use. For other uses you need to obtain permission from the rights-holder(s) directly, unless additional rights are indicated by a Creative Commons license in the record and/ or on the work itself. This material has been accepted for inclusion by faculty at Illinois Wesleyan University. For more information, please contact digitalcommons@iwu.edu.

©Copyright is owned by the author of this document.

β -AMYLOID (1-42) AND ITS ROLE IN DEVELOPING AN ANIMAL MODEL OF ALZHEIMER'S DISEASE

Greg Tinkler, and Dr. Wayne Dornan*,
Department of Psychology, IWU.

Alzheimer's disease is a neurodegenerative disorder characterized by neurofibrillary tangles and neuritic plaques. The major component of neuritic plaques is β -amyloid. In humans, β -amyloid is a chain of amino acids varying in length from 39 to 43. Current studies have shown that β -amyloid (1-40) is toxic to cells *in vitro*, with the 25-35 chain of amino acids supposedly being responsible for the cell death. However, the ability of the protein to cause cell death *in vivo* is still being questioned. One problem is that the (1-40) form has been found to aggregate very rapidly in most solvents, which could prevent a successful injection of β A into a rat. Recently, the (1-42) form has been purified and synthesized. As opposed to the (1-40) form, the new 42 amino acid sequence has a slower rate of sedimentation, which increases the likelihood of an effective injection. In order to determine the effect of β A (1-42) on spatial tasks, 0.5 microliters of the peptide were injected bilaterally into the dorsal hippocampus of male Long Evans rats. Control groups were also utilized. The first control group consisted of injections of a scrambled version of the peptide, the second group being the vehicle (DMSO) alone. The animals, which had been pretrained on a radial arm maze, were allowed to recuperate from surgery and were then tested on the maze again. The maze consisted of eight arms, five of which were always baited for a particular animal. Session latency, latency to first choice, number of correct choices, number of reference memory errors, number of correct and incorrect errors, and the total number of choices were recorded for each animal. Items that were then calculated included the percent of correct choices, the average choice latency, and the total number of errors. Testing continued for two weeks, after which the reinforcers from three of the arms were removed and shifted them to the three arms which had previously been unbaited. Three days later, all reinforcers were removed except for the two arms which had been baited for both of the previous configurations. These two reconfiguration procedures were done in an attempt to differentiate between animals using procedural memory and those relying on declarative memory, which is controlled by the hippocampus. The data was analyzed using ANOVA. The rats were perfused after testing, and the histologies are currently being completed. There was no significant difference between the three groups. These results suggest that injections of β A do not affect the retention of previously learned spatial tasks in the rat. Further studies will assess the affects of bilateral injections of β A (1-42) into the hippocampus on the acquisition of a spatial learning task.