



**Illinois Wesleyan University**  
**Digital Commons @ IWU**

---

John Wesley Powell Student Research  
Conference

1995, 6th Annual JWP Conference

---

Apr 22nd, 10:00 AM - 4:00 PM

## Nest Predation on House Wrens

Daniel L. Moore  
*Illinois Wesleyan University*

R. Given Harper, Faculty Advisor  
*Illinois Wesleyan University*

Follow this and additional works at: <https://digitalcommons.iwu.edu/jwprc>

---

Moore, Daniel L. and Harper, Faculty Advisor, R. Given, "Nest Predation on House Wrens" (1995). *John Wesley Powell Student Research Conference*. 16.  
<https://digitalcommons.iwu.edu/jwprc/1995/posters/16>

This Event is protected by copyright and/or related rights. It has been brought to you by Digital Commons @ IWU with permission from the rights-holder(s). You are free to use this material in any way that is permitted by the copyright and related rights legislation that applies to your use. For other uses you need to obtain permission from the rights-holder(s) directly, unless additional rights are indicated by a Creative Commons license in the record and/ or on the work itself. This material has been accepted for inclusion by faculty at Illinois Wesleyan University. For more information, please contact [digitalcommons@iwu.edu](mailto:digitalcommons@iwu.edu).

©Copyright is owned by the author of this document.

## NEST PREDATION ON HOUSE WRENS

Daniel L. Moore and R. Given Harper\*, Department of Biology, IWU

Predation is the strongest selection pressure placed on most avian species. Birds can potentially manipulate several variables, such as nest-site selection, brood size, and clutch size, in an attempt to decrease the frequency of predation and increase their fitness. The main focus of this study is to determine the effect of habitat edges (i.e. edge effects) on nest predation in a population of House Wrens (*Troglodytes aedon*) which breeds in nest boxes. Although the House Wren is not an endangered species, the patterns of edge effects are increasingly important in the conservation of bird species whose populations are threatened by habitat fragmentation. This study includes data on nest predation involving 585 nest boxes from 1983-1994. Our results showed no significant edge effect, but we did find a significant variation in the effect of the type of edge (abrupt vs. tapering vs. riverine). We also found the major predators were raccoons (*Procyon lotor*) and other House Wrens. Predation by other House Wrens was significantly higher along abrupt edges than tapered and riverine edges. While raccoon predation was significantly higher along riverine edges than abrupt and tapered edges. We also found that predation rates did not significantly differ between brood sizes, clutch sizes, and upland vs. floodplain habitats. A possible reason we found no significant edge effect is that the study area may be too small, and, in effect, the whole study area is subject to the edge effect. This conclusion would mean that a larger forest than our 108 ha. study area would be needed to reduce the edge effect on woodland avian species. Since edge type is significant, as long as the edge transition was gradual between the forest and the other habitat, predation rates would likely be diminished.