



Apr 18th, 9:00 AM - 10:30 AM

## Synthesis of a New Family of Diamines: Components for Supramolecular Architectures

Jonathan T. Brockman  
*Illinois Wesleyan University*

Rebecca Roesner, Faculty Advisor  
*Illinois Wesleyan University*

Follow this and additional works at: <https://digitalcommons.iwu.edu/jwprc>

---

Brockman, Jonathan T. and Roesner, Faculty Advisor, Rebecca, "Synthesis of a New Family of Diamines: Components for Supramolecular Architectures" (1998). *John Wesley Powell Student Research Conference. 2.*

<https://digitalcommons.iwu.edu/jwprc/1998/posters/2>

This is protected by copyright and/or related rights. It has been brought to you by Digital Commons @ IWU with permission from the rights-holder(s). You are free to use this material in any way that is permitted by the copyright and related rights legislation that applies to your use. For other uses you need to obtain permission from the rights-holder(s) directly, unless additional rights are indicated by a Creative Commons license in the record and/ or on the work itself. This material has been accepted for inclusion by faculty at Illinois Wesleyan University. For more information, please contact [digitalcommons@iwu.edu](mailto:digitalcommons@iwu.edu).

©Copyright is owned by the author of this document.

## Poster Presentation 2

SYNTHESIS OF A NEW FAMILY OF DIAMINES:  
COMPONENTS FOR SUPRAMOLECULAR ARCHITECTURESJonathan T. Brockman and Rebecca Roesner\*  
Department of Chemistry, Illinois Wesleyan University

The practical limits of manufacturing smaller and smaller electronic components is fast approaching, and scientists have been exploring the use of molecular and macromolecular electronic devices. Supramolecular systems have been designed and demonstrated to function as molecular switches, molecular wires or photoelectric devices. One supramolecular structure is the rotaxane which is composed of a linear molecular string (linker) threaded through a macrocyclic molecular ring with bulky blocking groups attached to the ends of the linker to prevent dethreading. The current goal is the synthesis of a family of diamine linkers of various lengths that can be attached to the bulky blocking group. Synthesis is being achieved through the reaction of *p*-hydroxyacetanilide with dibromoalkanes in basic media, followed by basic hydrolysis.<sup>1</sup> An example with dibromobutane is shown in Figure 1.




Figure 1

<sup>1</sup>Bartulin, J.; Ramos, M.L.; Rivas, B.L. *Polymer Bulletin* 15, 405-409 (1986).