



Illinois Wesleyan University
Digital Commons @ IWU

John Wesley Powell Student Research
Conference

1998, 9th Annual JWP Conference

Apr 18th, 9:00 AM - 10:30 AM

**The Developmental Response to Varying Growth Rates in
Romalea Guttata, the Eastern Lubber Grasshopper**

Virginia L. Flanagin
Illinois Wesleyan University

Steven A. Juliano, Faculty Advisor
Illinois Wesleyan University

Follow this and additional works at: <https://digitalcommons.iwu.edu/jwprc>

Flanagin, Virginia L. and Juliano, Faculty Advisor, Steven A., "The Developmental Response to Varying Growth Rates in *Romalea Guttata*, the Eastern Lubber Grasshopper" (1998). *John Wesley Powell Student Research Conference*. 6.
<https://digitalcommons.iwu.edu/jwprc/1998/posters/6>

This Event is protected by copyright and/or related rights. It has been brought to you by Digital Commons @ IWU with permission from the rights-holder(s). You are free to use this material in any way that is permitted by the copyright and related rights legislation that applies to your use. For other uses you need to obtain permission from the rights-holder(s) directly, unless additional rights are indicated by a Creative Commons license in the record and/ or on the work itself. This material has been accepted for inclusion by faculty at Illinois Wesleyan University. For more information, please contact digitalcommons@iwu.edu.

©Copyright is owned by the author of this document.

Poster Presentation 6

**THE DEVELOPMENTAL RESPONSE TO VARYING GROWTH RATES
IN *ROMALEA GUTTATA*, THE EASTERN LUBBER GRASSHOPPER**

Virginia L. Flanagan and Steven A. Juliano*

Department of Biology, Illinois Wesleyan University and Illinois State University

Life history traits such as the size and relative age of an organism can greatly influence an individual's survival and reproductive success. These traits can vary between individuals in a species, due to environmental conditions during juvenile development. We examined how changes in food availability during juvenile stages of development can affect the size and timing of eclosion (i.e. adulthood) in *Romalea guttata*, the Eastern Lubber grasshopper. Hatchling *R. guttata* were raised on nine treatment groups consisting of a high food diet, a low food diet and switches from low to high food, high to low food, and high to no food diets. The switches were made at significant points in the insects development corresponding to different instars (i.e. stages of development). When food availability was high in the second through the fourth instars insects took significantly less time to reach adulthood than insects with low food availability in the middle three instars. The same trend occurred in the mass of the insects at eclosion; when high food availability occurred in the middle three instars, insects weighed significantly more than insects fed low food in the middle three instars. This suggests that both the time to and size at eclosion were determined in the middle of the nymphal life cycle, and that feeding in the first and last instar did not affect development. Other tests were done to examine whether insects with low food availability could increase their feeding efficiency, but our results did not support this hypothesis. Our results suggest that developmental cues are set at a particular developmental state, and proceed independently of feeding in the later instars. This suggests that development does not continuously adapt to changes in food availability, and there is strong evidence for biological cues which can cause development to proceed regardless of feeding. The concept of fixed or unresponsive development, in turn, may constrain an individual's ability to adapt to varying environments, resulting in a decrease in survival and reproductive success.