



Illinois Wesleyan University
Digital Commons @ IWU

John Wesley Powell Student Research
Conference

1998, 9th Annual JWP Conference

Apr 18th, 9:00 AM - 10:30 AM

Systematic Relationships of Rhinolophid Bats, Based on Hyoid Morphology

Jennifer L. Garner
Illinois Wesleyan University

Thomas A. Griffiths, Faculty Advisor
Illinois Wesleyan University

Follow this and additional works at: <https://digitalcommons.iwu.edu/jwprc>

Garner, Jennifer L. and Griffiths, Faculty Advisor, Thomas A., "Systematic Relationships of Rhinolophid Bats, Based on Hyoid Morphology" (1998). *John Wesley Powell Student Research Conference*. 7.

<https://digitalcommons.iwu.edu/jwprc/1998/posters/7>

This Event is protected by copyright and/or related rights. It has been brought to you by Digital Commons @ IWU with permission from the rights-holder(s). You are free to use this material in any way that is permitted by the copyright and related rights legislation that applies to your use. For other uses you need to obtain permission from the rights-holder(s) directly, unless additional rights are indicated by a Creative Commons license in the record and/ or on the work itself. This material has been accepted for inclusion by faculty at Illinois Wesleyan University. For more information, please contact digitalcommons@iwu.edu.

©Copyright is owned by the author of this document.

Poster Presentation 7

SYSTEMATIC RELATIONSHIPS OF RHINOLOPHID BATS, BASED ON
HYOID MORPHOLOGY

Jennifer L. Garner and Thomas A. Griffiths*
Department of Biology, Illinois Wesleyan University

Using standard microdissection techniques, the morphology of the hyoid apparatus and musculature was examined in two species of rhinolophid bats. A cladistic analysis of the data revealed that there were thirteen derived characters which could be used in the study. As expected, hyoid data support the traditional grouping of the family Rhinolophidae with the bat families Hipposideridae, Megadermatidae and Nycteridae. However, the results show, surprisingly, that these four families were on a line at the base of the microchiropteran tree, with all other bat families grouped in a clade above them. This study is the first to suggest that the rhinolophid clade is the basal line of the microchiropteran tree.