



Apr 18th, 9:00 AM - 10:30 AM

## Synthesis of a Fully Conjugated Linear Diamine as a Potential Rotaxane Component

Matthew T. Lewellen  
*Illinois Wesleyan University*

Rebecca Roesner, Faculty Advisor  
*Illinois Wesleyan University*

Follow this and additional works at: <https://digitalcommons.iwu.edu/jwprc>

---

Lewellen, Matthew T. and Roesner, Faculty Advisor, Rebecca, "Synthesis of a Fully Conjugated Linear Diamine as a Potential Rotaxane Component" (1998). *John Wesley Powell Student Research Conference*. 13.  
<https://digitalcommons.iwu.edu/jwprc/1998/posters/13>

This is protected by copyright and/or related rights. It has been brought to you by Digital Commons @ IWU with permission from the rights-holder(s). You are free to use this material in any way that is permitted by the copyright and related rights legislation that applies to your use. For other uses you need to obtain permission from the rights-holder(s) directly, unless additional rights are indicated by a Creative Commons license in the record and/ or on the work itself. This material has been accepted for inclusion by faculty at Illinois Wesleyan University. For more information, please contact [digitalcommons@iwu.edu](mailto:digitalcommons@iwu.edu).

©Copyright is owned by the author of this document.

Poster Presentation 13

SYNTHESIS OF A FULLY CONJUGATED LINEAR DIAMINE AS A POTENTIAL ROTAXANE COMPONENT

Matthew T. Lewellen and Rebecca A. Roesner\*  
 Department of Chemistry, Illinois Wesleyan University

Steps toward the preparation of a fully conjugated linear diamine are described. This diamine will be used as a linear component in the preparation of larger supramolecular species called rotaxanes. Rotaxanes are composed of a ring shaped molecule threaded onto a linear molecule with bulky stoppering groups used to prevent dethreading of the ring (Figure 1). The target rotaxane will be composed of two hexamolybdate ions, a beta-cyclodextrin molecule, and a linear diamine. In the first step of the diamine synthesis, *p*-aminotrimethylsilylethynylbenzene was prepared from 4-iodoaniline and trimethylsilyl acetylene according to Figure 2. The trimethylsilyl protecting group allows substitution to occur at only one C-H terminus of the acetylene molecule. In a subsequent step (Figure 3) the trimethylsilyl protecting group was removed from the *p*-aminotrimethylsilylethynylbenzene by addition of aqueous potassium hydroxide. In a third step, two equivalents of the newly formed *p*-aminoethynylbenzene will be coupled together to form 4,4'-bis(amino)diphenylbutadiyne (Figure 4).




Figure 1




Figure 2




Figure 3




Figure 4