



**Illinois Wesleyan University**  
**Digital Commons @ IWU**

---

John Wesley Powell Student Research  
Conference

1998, 9th Annual JWP Conference

---

Apr 18th, 9:00 AM - 10:30 AM

## Substitution Chemistry of the Hexamolybdate Ion: The Development of Reactions for Rotaxane Preparation

Sara C. McGrath  
*Illinois Wesleyan University*

Rebecca A. Roesner, Faculty Advisor  
*Illinois Wesleyan University*

Follow this and additional works at: <https://digitalcommons.iwu.edu/jwprc>

---

McGrath, Sara C. and Roesner, Faculty Advisor, Rebecca A., "Substitution Chemistry of the Hexamolybdate Ion: The Development of Reactions for Rotaxane Preparation" (1998). *John Wesley Powell Student Research Conference*. 14.  
<https://digitalcommons.iwu.edu/jwprc/1998/posters/14>

This Event is protected by copyright and/or related rights. It has been brought to you by Digital Commons @ IWU with permission from the rights-holder(s). You are free to use this material in any way that is permitted by the copyright and related rights legislation that applies to your use. For other uses you need to obtain permission from the rights-holder(s) directly, unless additional rights are indicated by a Creative Commons license in the record and/ or on the work itself. This material has been accepted for inclusion by faculty at Illinois Wesleyan University. For more information, please contact [digitalcommons@iwu.edu](mailto:digitalcommons@iwu.edu).

©Copyright is owned by the author of this document.

## Poster Presentation 14

## SUBSTITUTION CHEMISTRY OF THE HEXAMOLYBDATE ION: THE DEVELOPMENT OF REACTIONS FOR ROTAXANE PREPARATION

Sara C. McGrath and Rebecca A. Roesner\*  
 Department of Chemistry, Illinois Wesleyan University

Much interest has been generated lately in the synthesis of new materials to be used in the construction of molecular scale electronic devices and machinery. Rotaxanes have surfaced as potential building blocks for designing such structures. The simplest rotaxanes are dumbbell-shaped molecules consisting of a long chain linker unit threaded, non-covalently, through a ring-shaped unit and stoppered on each end with a large, bulky group. (Fig. 1) The object of this study is to synthesize a simple rotaxane, using various bifunctional organic molecules as the chain unit, a  $\beta$ -cyclodextrin molecule as the ring unit, and hexamolybdate ions as the stoppering groups. IR and  $^1\text{H}$  NMR analyses indicate that the hexamolybdate ion has been successfully synthesized from  $\text{Na}_2\text{MoO}_4$  and  $(n\text{-C}_4\text{H}_9)_4\text{NBr}$  (Inorganic Syntheses). The main focus of this project is to explore the substitution reactions between the hexamolybdate ion and various aromatic amines, and, by analogy, to understand the chemistry of the substitution reactions between the hexamolybdate ion and the potential rotaxane linkers. There has been some preliminary success with the substitution of *p*-anisidine and 1,4-phenylenediamine onto the hexamolybdate ion, using high pressure sealed tube reactions at elevated temperatures. Analyses performed using  $^1\text{H}$  NMR and FAB-MS suggest that the aromatic molecules have been successfully substituted onto the hexamolybdate ion. (Fig. 2)




Fig. 1




Fig. 2