



**Illinois Wesleyan University**  
**Digital Commons @ IWU**

---

John Wesley Powell Student Research  
Conference

1998, 9th Annual JWP Conference

---

Apr 18th, 1:30 PM - 2:30 PM

## **The Role of the Comstock-Kellogg Glands in Egg Tanning in *Romalea Guttata***

Elizabeth Elsasser  
*Illinois Wesleyan University*

Given Harper, Faculty Advisor  
*Illinois Wesleyan University*

Follow this and additional works at: <https://digitalcommons.iwu.edu/jwprc>

---

Elsasser, Elizabeth and Harper, Faculty Advisor, Given, "The Role of the Comstock-Kellogg Glands in Egg Tanning in *Romalea Guttata*" (1998). *John Wesley Powell Student Research Conference*. 6.

<https://digitalcommons.iwu.edu/jwprc/1998/posters2/6>

This Event is protected by copyright and/or related rights. It has been brought to you by Digital Commons @ IWU with permission from the rights-holder(s). You are free to use this material in any way that is permitted by the copyright and related rights legislation that applies to your use. For other uses you need to obtain permission from the rights-holder(s) directly, unless additional rights are indicated by a Creative Commons license in the record and/ or on the work itself. This material has been accepted for inclusion by faculty at Illinois Wesleyan University. For more information, please contact [digitalcommons@iwu.edu](mailto:digitalcommons@iwu.edu).

©Copyright is owned by the author of this document.

Poster Presentation 24

THE ROLE OF THE COMSTOCK-KELLOGG GLANDS IN EGG  
TANNING IN *ROMALEA GUTTATA*

Elizabeth Elsasser and Given Harper\*

Department of Biology, Illinois Wesleyan University

Freshly laid eggs of the Eastern Lubber grasshopper, *Romalea guttata*, are yellow and soft-shelled, but within several hours the eggs become hardened and dark brown. In a similar process, calyx and oviduct secretions, which form the egg pod, become frothy, darkened, hardened, and water insoluble during oviposition. It has been suggested that secretions from the paired Comstock-Kellogg glands accelerate tanning of both the eggs and egg pod foam. I investigated the effects of Comstock-Kellogg gland secretions on the rate of egg and egg pod tanning during egg pod production in the female Eastern Lubber grasshopper. Eggs streaked with CK gland secretions, macerated CK gland, macerated intersegmental membrane, or hemolymph took significantly less time to tan than eggs streaked with water or nothing. Females lacking CK glands laid eggs that took significantly longer to tan than eggs from sham-operated or unoperated females. These results suggest that the CK glands do serve to accelerate tanning, but the chemical composition of the CK secretions may be common to many grasshopper tissues.