



Apr 15th, 11:00 AM - 12:30 PM

## **Brown-Headed Cowbird Brood Parasitism in Bison-Grazed and Ungrazed Tall Grass Prairie in Northeastern Kansas**

Alexandra S. Latham  
*Illinois Wesleyan University*

Jack F. Cully, Faculty Advisor  
*Kansas State University*

Sheryl Swartz Soukup, Faculty Advisor  
*Illinois Wesleyan University*

Follow this and additional works at: <https://digitalcommons.iwu.edu/jwprc>

---

Latham, Alexandra S.; Cully, Faculty Advisor, Jack F.; and Soukup, Faculty Advisor, Sheryl Swartz, "Brown-Headed Cowbird Brood Parasitism in Bison-Grazed and Ungrazed Tall Grass Prairie in Northeastern Kansas" (2000). *John Wesley Powell Student Research Conference*. 3.

<https://digitalcommons.iwu.edu/jwprc/2000/oralpres3/3>

This Event is protected by copyright and/or related rights. It has been brought to you by Digital Commons @ IWU with permission from the rights-holder(s). You are free to use this material in any way that is permitted by the copyright and related rights legislation that applies to your use. For other uses you need to obtain permission from the rights-holder(s) directly, unless additional rights are indicated by a Creative Commons license in the record and/ or on the work itself. This material has been accepted for inclusion by faculty at Illinois Wesleyan University. For more information, please contact [digitalcommons@iwu.edu](mailto:digitalcommons@iwu.edu).

©Copyright is owned by the author of this document.

Oral Presentation 3.3  
(Also see Poster 28)

**BROWN -HEADED COWBIRD BROOD PARASITISM IN BISON-GRAZED AND  
UNGRAZED TALLGRASS PRAIRIE IN NORTHEASTERN KANSAS**

Alexandra S. Latham and Jack F. Cully\*, Department of Biological Sciences, Kansas  
State University, and Sheryl Swartz Soukup\*, Department of Biology,  
Illinois Wesleyan University

The Brown-headed Cowbird (*Molothrus ater*) has long been associated with bison (*Bos bison*) in North America on the Great Plains. As a result, we anticipated that cowbirds would be more successful breeding in the presence of bison than in their absence. We predicted that several common ground-nesting avian species, Dickcissels (*Spiza americana*), Grasshopper Sparrows (*Ammodramus savannarum*), and Eastern Meadowlarks (*Sturnella magna*), would suffer higher frequencies of brood parasitism in bison-grazed habitat than in ungrazed habitat on Konza Prairie Research Natural Area in northeastern Kansas. The frequency of cowbird parasitism for all species combined was significantly higher (0.69) in bison-grazed than in ungrazed habitat (0.44) ( $p = 0.044$ ,  $\chi^2 = 4.061$ ,  $df = 1$ ). These results are consistent with our suggestion that bison-grazed habitat may be a more optimal site for cowbird brood parasitism than ungrazed habitat. We pose two principal explanations for the higher frequency of parasitism observed in the bison-grazed area. First, cowbirds may be able to forage more efficiently in the bison-grazed area, indirectly inflating parasitism frequencies by conferring a variety of energetic and nutritional advantages upon the females. Second, the cowbirds' abilities to find and parasitize nests may be enhanced by the shorter, less dense grass characteristic of grazed habitat. Further studies investigating the conservation implications of this phenomenon are merited since cowbird brood parasitism usually reduces reproductive success of host species, and has had a dramatic negative impact on population of several hosts, driving them to near extinction.