



Apr 12th, 9:00 AM - 10:30 AM

## Developing an Animal Model of Alzheimer's Disease: Multiple Hippocampal Injections of Beta-Amyloid (1-42) and the Effects on Spatial Learning and Memory in the Male Rat

Gregory P. Tinkler  
*Illinois Wesleyan University*

Wayne A. Dornan, Faculty Advisor  
*Illinois Wesleyan University*

Follow this and additional works at: <https://digitalcommons.iwu.edu/jwprc>

---

Tinkler, Gregory P. and Dornan, Faculty Advisor, Wayne A., "Developing an Animal Model of Alzheimer's Disease: Multiple Hippocampal Injections of Beta-Amyloid (1-42) and the Effects on Spatial Learning and Memory in the Male Rat" (1997). *John Wesley Powell Student Research Conference*. 25.

<https://digitalcommons.iwu.edu/jwprc/1997/posters/25>

This Event is protected by copyright and/or related rights. It has been brought to you by Digital Commons @ IWU with permission from the rights-holder(s). You are free to use this material in any way that is permitted by the copyright and related rights legislation that applies to your use. For other uses you need to obtain permission from the rights-holder(s) directly, unless additional rights are indicated by a Creative Commons license in the record and/ or on the work itself. This material has been accepted for inclusion by faculty at Illinois Wesleyan University. For more information, please contact [digitalcommons@iwu.edu](mailto:digitalcommons@iwu.edu).

©Copyright is owned by the author of this document.

Poster Presentation 23

**DEVELOPING AN ANIMAL MODEL OF ALZHEIMER'S DISEASE:  
MULTIPLE HIPPOCAMPAL INJECTIONS OF BETA-AMYLOID (1-42)  
AND THE EFFECTS ON SPATIAL LEARNING AND MEMORY IN  
THE MALE RAT.**

Gregory P. Tinkler and Wayne A. Dornan\*, Department of Psychology, IWU

Alzheimer's Disease (AD) is a neurodegenerative disorder which affects approximately four million people in the United States alone. A diagnosis of AD can be made only by performing a postmortem examination of the brain and confirming the presence of high numbers of neuritic plaques and neurofibrillary tangles, the pathological hallmarks of the disease. The neuritic plaque is composed primarily of the 39- to 43- amino acid protein  $\beta$ -Amyloid ( $\beta$ A), and is found in large quantities in structures such as the hippocampus, which is intimately involved with learning and memory.  $\beta$ A has been hypothesized to play a role in the onset of AD, but studies which attempt to model the disease using single injections of the peptide fail to produce consistent behavioral impairments. A number of recent studies have suggested that chronic administration of  $\beta$ A or one of its fragments may be more pathologically relevant to AD than single injections, and may also produce more consistent impairments of spatial learning and memory. In addition, our lab has demonstrated that multiple injections of the neurotoxic fragment  $\beta$ A (25-35) into the septum, which projects to the hippocampus, produce a marginal impairment on a spatial task. A marginal depletion of cholinergic markers in the hippocampus was also produced. Since the  $\beta$ A fragment has different chemical properties than the full length peptide, it is conceivable that more profound behavioral and neurochemical impairments may result from the use of  $\beta$ A (1-42). It is also possible that injections into the hippocampus instead of the septum would produce a greater impairment on learning and memory. Therefore, this study examines the effects of multiple hippocampal injections of the full length  $\beta$ A (1-42) peptide on spatial learning and memory in the rat.