



Illinois Wesleyan University
Digital Commons @ IWU

John Wesley Powell Student Research
Conference

1997, 8th Annual JWP Conference

Apr 12th, 9:00 AM - 10:30 AM

Development of a Simple, Inexpensive, and Rapid Screening Method for Detection of Cocaine Use

Jennifer A. Duke
Illinois Wesleyan University

David N. Bailey, Faculty Advisor
Illinois Wesleyan University

Follow this and additional works at: <https://digitalcommons.iwu.edu/jwprc>

Duke, Jennifer A. and Bailey, Faculty Advisor, David N., "Development of a Simple, Inexpensive, and Rapid Screening Method for Detection of Cocaine Use" (1997). *John Wesley Powell Student Research Conference*. 32.
<https://digitalcommons.iwu.edu/jwprc/1997/posters/32>

This Event is protected by copyright and/or related rights. It has been brought to you by Digital Commons @ IWU with permission from the rights-holder(s). You are free to use this material in any way that is permitted by the copyright and related rights legislation that applies to your use. For other uses you need to obtain permission from the rights-holder(s) directly, unless additional rights are indicated by a Creative Commons license in the record and/ or on the work itself. This material has been accepted for inclusion by faculty at Illinois Wesleyan University. For more information, please contact digitalcommons@iwu.edu.

©Copyright is owned by the author of this document.

Poster Presentation 31

**DEVELOPMENT OF A SIMPLE, INEXPENSIVE, AND RAPID
SCREENING METHOD FOR DETECTION OF COCAINE USE**

Jennifer A. Duke and David N. Bailey*, Department of Chemistry, IWU

A quick, inexpensive, and accurate method of analyzing urine to detect drug use is needed for the handling of large numbers of samples commonly encountered in screening applications. One method currently under investigation for the detection of cocaine derivatives in human urine is ion-ion pairing. A version of this technique, utilizing tetrakis (thiocyanato) cobalt II as the pairing agent, has been applied to facilitate extraction of the cocaine metabolite benzoylecgonine (BE) into an organic solvent. The efficiency of this procedure is dependent upon pH and ionic strength of the solution. The most favorable conditions for the extraction of BE into dichloromethane have been determined.