



Apr 15th, 2:00 PM - 3:00 PM

## Attaching Organic Ligands to the Keggin Ion

Jacqui Hoffman

*Illinois Wesleyan University*

Rebecca Roesner, Faculty Advisor

*Illinois Wesleyan University*

Follow this and additional works at: <http://digitalcommons.iwu.edu/jwprc>

---

Hoffman, Jacqui and Roesner, Faculty Advisor, Rebecca, "Attaching Organic Ligands to the Keggin Ion" (2000). *John Wesley Powell Student Research Conference*. 21.

<http://digitalcommons.iwu.edu/jwprc/2000/posters2/21>

This Event is brought to you for free and open access by The Ames Library, the Andrew W. Mellon Center for Curricular and Faculty Development, the Office of the Provost and the Office of the President. It has been accepted for inclusion in Digital Commons @ IWU by the faculty at Illinois Wesleyan University. For more information, please contact [digitalcommons@iwu.edu](mailto:digitalcommons@iwu.edu).

©Copyright is owned by the author of this document.

Poster Presentation 18

**ATTACHING ORGANIC LIGANDS TO THE KEGGIN ION**

Jacqui Hoffman and Rebecca Roesner\*  
Department of Chemistry, Illinois Wesleyan University

Polyoxometalates are early transition metal-oxygen cluster compounds. Such complexes are known to be valuable catalysts, analytical reagents, and electron microscopy stains. Another potentially important application of heteropoly species is in clinical medicine. Several polytungstates have significant antiviral and antitumoral activity. The Keggin anion,  $\text{XM}_{12}\text{O}_{40}^{-n}$ , shown in figure 1, is among the most biologically active polyoxometalates.

By attaching organic molecules to the surface of this cluster, it may be possible to




Figure 1

develop drugs which "seek out" diseased cells. Before this can be accomplished, however, fundamental techniques for forming polyoxometalate-carbon bonds must be developed.

The first molecule that we synthesized was a lacunary Keggin ion. A lacunary ion has one W-O group missing resulting in a vacancy in an otherwise symmetric molecule. The structure was confirmed by  $^{31}\text{P}$ NMR, FT-IR and UV analysis. A carboxylate functional group was then attached to the lacunary ion by inserting a rhodium atom bearing a  $-\text{CH}_2\text{COOH}$  ligand. The color change from white to orange (indicative of rhodium) signified successful substitution of the vacancy. The product was analyzed by  $^1\text{H}$ NMR and FT-IR. A  $^{31}\text{P}$ NMR spectrum was also obtained which showed the expected chemical shift. In the next and final step, we attempted to convert the carboxylate group to an amide through an anhydride intermediate.