



Apr 12th, 1:30 PM - 2:30 PM

## Phylogenetic Systematics of the Primate Genus *Aotus* Based on Hyoid Morphology

Melissa S. Immel  
*Illinois Wesleyan University*

Thomas A. Griffiths, Faculty Advisor  
*Illinois Wesleyan University*

Follow this and additional works at: <https://digitalcommons.iwu.edu/jwprc>

---

Immel, Melissa S. and Griffiths, Faculty Advisor, Thomas A., "Phylogenetic Systematics of the Primate Genus *Aotus* Based on Hyoid Morphology" (1997). *John Wesley Powell Student Research Conference*. 37.

<https://digitalcommons.iwu.edu/jwprc/1997/posters/37>

This Event is protected by copyright and/or related rights. It has been brought to you by Digital Commons @ IWU with permission from the rights-holder(s). You are free to use this material in any way that is permitted by the copyright and related rights legislation that applies to your use. For other uses you need to obtain permission from the rights-holder(s) directly, unless additional rights are indicated by a Creative Commons license in the record and/ or on the work itself. This material has been accepted for inclusion by faculty at Illinois Wesleyan University. For more information, please contact [digitalcommons@iwu.edu](mailto:digitalcommons@iwu.edu).

©Copyright is owned by the author of this document.

Poster Presentation 38

**PHYLOGENETIC SYSTEMATICS OF THE PRIMATE GENUS *AOTUS*,  
BASED ON HYOID MORPHOLOGY**

Melissa S. Immel and Thomas A. Griffiths\*, Department of Biology, IWU

Dissection of a specimen of the South American primate genus Aotus was performed, focussing on the morphology of the hyoid apparatus and surrounding musculature. Data collected from this dissection, along with data from published descriptions of primate genera from both suborders Haplorhini and Strepsirhini, were analyzed using the computer software program PAUP (Phylogenetic Analysis Using Parsimony). This analysis generates possible phylogenetic relationships of Aotus to other primates. Four equally probable phylogenetic trees (=cladograms) were obtained from this analysis. While the cladograms differ in minor detail, they all show that Aotus appears to be more closely related to the more advanced suborder Haplorhini than to the Strepsirhini, and that it is a relatively primitive member of the haplorhine group. This suggests that the ancestor of the New World monkeys arrived in South America very early, at about the time that haplorhine primates were beginning to diverge in the Old World.