



**Illinois Wesleyan University**  
**Digital Commons @ IWU**

---

John Wesley Powell Student Research  
Conference

2000, 11th Annual JWP Conference

---

Apr 15th, 10:00 AM - 11:00 AM

## Identification of LRF1 and LRF2 Mutants in Arabidopsis

Neva Laurie  
*Illinois Wesleyan University*

Loni Walker, Faculty Advisor  
*Illinois Wesleyan University*

Follow this and additional works at: <https://digitalcommons.iwu.edu/jwprc>

---

Laurie, Neva and Walker, Faculty Advisor, Loni, "Identification of LRF1 and LRF2 Mutants in Arabidopsis" (2000). *John Wesley Powell Student Research Conference*. 14.  
<https://digitalcommons.iwu.edu/jwprc/2000/posters/14>

This Event is protected by copyright and/or related rights. It has been brought to you by Digital Commons @ IWU with permission from the rights-holder(s). You are free to use this material in any way that is permitted by the copyright and related rights legislation that applies to your use. For other uses you need to obtain permission from the rights-holder(s) directly, unless additional rights are indicated by a Creative Commons license in the record and/ or on the work itself. This material has been accepted for inclusion by faculty at Illinois Wesleyan University. For more information, please contact [digitalcommons@iwu.edu](mailto:digitalcommons@iwu.edu).

©Copyright is owned by the author of this document.

Poster Presentation 29

**IDENTIFICATION OF LRF1 AND LRF2 MUTANTS IN ARABIDOPSIS**

Neva Laurie and Loni Walker\*

Department of Biology, Illinois Wesleyan University

Arabidopsis is a model system for research in plant genetics. It is a typical dicot with a relatively short life cycle and has a small, compact genome with little noncoding DNA. Because these characteristics make it so appealing for research, much of its genome has been mapped, resulting in a database of known genes with unknown functions. The purpose of this experiment is to determine a function for two of these genes, LRF1 and LRF2. These genes are suspected to code for F-box proteins, which are believed to provide specificity in marking proteins for degradation through the ubiquitin pathway. This function often acts as a control mechanism in hormone signaling pathways, and sequence similarity to known genes suggests this is a likely mode of action for LRF1 and LRF2. This experiment employs reverse genetic techniques to identify plants which have had these genes mutated, causing production of a nonfunctional protein.