



Apr 15th, 2:00 PM - 3:00 PM

## Synthesis of Bis(Hexamolybdate) Complexes Covalently Bridged by Organnoimido Ligands

Jeremie D. Moll  
*Illinois Wesleyan University*

Rebecca Roesner, Faculty Advisor  
*Illinois Wesleyan University*

Follow this and additional works at: <https://digitalcommons.iwu.edu/jwprc>

---

Moll, Jeremie D. and Roesner, Faculty Advisor, Rebecca, "Synthesis of Bis(Hexamolybdate) Complexes Covalently Bridged by Organnoimido Ligands" (2000). *John Wesley Powell Student Research Conference*. 15.  
<https://digitalcommons.iwu.edu/jwprc/2000/posters2/15>

This is protected by copyright and/or related rights. It has been brought to you by Digital Commons @ IWU with permission from the rights-holder(s). You are free to use this material in any way that is permitted by the copyright and related rights legislation that applies to your use. For other uses you need to obtain permission from the rights-holder(s) directly, unless additional rights are indicated by a Creative Commons license in the record and/ or on the work itself. This material has been accepted for inclusion by faculty at Illinois Wesleyan University. For more information, please contact [digitalcommons@iwu.edu](mailto:digitalcommons@iwu.edu).

©Copyright is owned by the author of this document.

Poster Presentation 34

SYNTHESIS OF BIS(HEXAMOLYBDATE) COMPLEXES  
COVALENTLY BRIDGED BY ORGANOIMIDO LIGANDS

Jeremie D. Moll and Rebecca Roesner\*  
Department of Chemistry, Illinois Wesleyan University




Figure 1

Rotaxanes are supermolecular structures composed of linear molecules threaded through macrocyclic molecular rings and held in place by the addition of large blocking groups to prevent dethreading (**Figure 1**). Polyoxometalates, such as hexamolybdate ions, are extremely large, highly symmetric clusters that can be used as the blocking groups for rotaxane preparation. Our intent has been to synthesize a Bis(Hexamolybdate) complex to serve as the dumbbell for our rotaxane through the use of difunctional amine linkers. Our current goal is to purify the synthesized complex (**Figure 2**) and then repeat the procedure with the macrocyclic molecule in place.

Figure 2

