

Illinois Wesleyan University Digital Commons @ IWU

John Wesley Powell Student Research Conference

1996, 7th Annual JWP Conference

Apr 13th, 1:30 PM - 2:30 PM

## Bacteria in House Wren (Troglodytes aedon) Nests

David R. Singleton Illinois Wesleyan University

R. Given Harper, Faculty Advisor *Illinois Wesleyan University* 

Follow this and additional works at: https://digitalcommons.iwu.edu/jwprc

Singleton, David R. and Harper, Faculty Advisor, R. Given, "Bacteria in House Wren (*Troglodytes aedon*) Nests" (1996). *John Wesley Powell Student Research Conference*. 11. https://digitalcommons.iwu.edu/jwprc/1996/posters/11

This Event is protected by copyright and/or related rights. It has been brought to you by Digital Commons @ IWU with permission from the rights-holder(s). You are free to use this material in any way that is permitted by the copyright and related rights legislation that applies to your use. For other uses you need to obtain permission from the rights-holder(s) directly, unless additional rights are indicated by a Creative Commons license in the record and/ or on the work itself. This material has been accepted for inclusion by faculty at Illinois Wesleyan University. For more information, please contact digitalcommons@iwu.edu.

 $\ensuremath{\mathbb{C}}$  Copyright is owned by the author of this document.

## Poster Presentation 34

## BACTERIA IN HOUSE WREN (Troglodytes aedon) NESTS

## David R. Singleton and R. Given Harper\*, Department of Biology, IWU

Nest ectoparasites such as mites, fleas, fly larvae, etc., have been studied extensively as selective agents in the evolution of life history traits in several species of birds. The potential influence of bacterial populations in nests, however, has not been investigated. In this study we are examining used House Wren nests to document the presence of bacterial species, some of which may be potential pathogens. Standard microbial techniques are currently being utilized to identify bacteria isolated from the nests. The presence of pathogens in the nest may help explain the particular behavior of male House Wrens and other cavity nesting birds in removing old nests from nesting sites before beginning the construction of new nests.