



Apr 12th, 9:00 AM - 10:00 AM

## Synthesis of Hexamolybdate-Polyether Diamine Complexes

David Aggen  
*Illinois Wesleyan University*

Rebecca Roesner, Faculty Advisor  
*Illinois Wesleyan University*

Follow this and additional works at: <https://digitalcommons.iwu.edu/jwprc>

---

Aggen, David and Roesner, Faculty Advisor, Rebecca, "Synthesis of Hexamolybdate-Polyether Diamine Complexes" (2003). *John Wesley Powell Student Research Conference*.  
1.

<https://digitalcommons.iwu.edu/jwprc/2003/posters/1>

This is protected by copyright and/or related rights. It has been brought to you by Digital Commons @ IWU with permission from the rights-holder(s). You are free to use this material in any way that is permitted by the copyright and related rights legislation that applies to your use. For other uses you need to obtain permission from the rights-holder(s) directly, unless additional rights are indicated by a Creative Commons license in the record and/ or on the work itself. This material has been accepted for inclusion by faculty at Illinois Wesleyan University. For more information, please contact [digitalcommons@iwu.edu](mailto:digitalcommons@iwu.edu).

©Copyright is owned by the author of this document.


## Poster Presentation P1

## SYNTHESIS OF HEXAMOLYBDATE-POLYETHER DIAMINE COMPLEXES

David Aggen and Rebecca Roesner\*

Department of Chemistry, Illinois Wesleyan University

Polyoxometalates are a class of highly symmetric anion clusters that have applications in supramolecular chemistry. These polyoxometalate clusters may be able to act as “ends” which trap macrocyclic rings on linear carbon chains. Polyoxometalates have already been incorporated into polymers<sup>1</sup> and dendrimers.<sup>2</sup> Our research involves the synthesis of a dumbbell shaped molecule through reaction of two equivalents of tetrabutylammonium hexamolybdate ( $[\text{Bu}_4\text{N}]_4[\text{Mo}_6\text{O}_{19}]$ ) and 1 equivalent of a polyether diamine strap. The diamine was prepared from 4-acetamidophenol and 1,2 bis(2-chloroethoxy)ethane. Further reaction of the prepared diamine with the hexamolybdate ion  $[\text{Mo}_6\text{O}_{19}]^{2-}$  was facilitated using dicyclohexylcarbodiimide ( $\text{C}_{13}\text{H}_{22}\text{N}_2$ ) which acts as a catalyst and a dehydrating agent. Dicyclohexylurea ( $\text{C}_{13}\text{H}_{24}\text{N}_2\text{O}$ ) is formed as a byproduct.<sup>3</sup> Formation of the diamine-hexamolybdate adduct was verified by  $^1\text{H}$  NMR and FT IR. We hope to refine our synthetic procedure to achieve higher yields.



<sup>1</sup> Moore, Aaron R.; Kwen, Haidoo; Beatty, Alicia M.; Maata, Eric A. *Chem. Comm.* **2000**, 1793-1794.

<sup>2</sup> Zeng, Huadong; Newkome, George R.; Hill, Craig L. *Agnew. Chem. Int. Ed.* **2000**, 39, No. 10, 1772-1774.