



Illinois Wesleyan University
Digital Commons @ IWU

John Wesley Powell Student Research
Conference

2003, 14th Annual JWP Conference

Apr 12th, 9:00 AM - 10:00 AM

Linking Polyoxometalates With Amide Bonds

Joseph B. Binder
Illinois Wesleyan University

Emilia Poppe
Illinois Wesleyan University

Rebecca A. Roesner, Faculty Advisor
Illinois Wesleyan University

Follow this and additional works at: <https://digitalcommons.iwu.edu/jwprc>

Binder, Joseph B.; Poppe, Emilia; and Roesner, Faculty Advisor, Rebecca A., "Linking Polyoxometalates With Amide Bonds" (2003). *John Wesley Powell Student Research Conference*. 7.

<https://digitalcommons.iwu.edu/jwprc/2003/posters/7>

This Event is protected by copyright and/or related rights. It has been brought to you by Digital Commons @ IWU with permission from the rights-holder(s). You are free to use this material in any way that is permitted by the copyright and related rights legislation that applies to your use. For other uses you need to obtain permission from the rights-holder(s) directly, unless additional rights are indicated by a Creative Commons license in the record and/ or on the work itself. This material has been accepted for inclusion by faculty at Illinois Wesleyan University. For more information, please contact digitalcommons@iwu.edu.

©Copyright is owned by the author of this document.


Poster Presentation P7

LINKING POLYOXOMETALATES WITH AMIDE BONDS

Joseph B. Binder, Emilia Poppe, and Rebecca A. Roesner*
Department of Chemistry, Illinois Wesleyan University

Polyoxometalates (POMs) of the Keggin structure, $[XM_{12}O_{40}]^{n-}$, are large metal-oxygen anions. One type, the unadecatungstophosphate lacunary Keggin ion ($[PW_{11}O_{39}]^{7-}$) has one fewer W-O unit than the usual Keggin ion, leaving a hole into which another metal atom can be inserted. In previous work, we inserted a rhodium atom with a carboxylate ligand into the vacancy and reacted this product with aniline to form an amide.

In the current work, the amide-forming reaction has been attempted with a diamine ($H_2NPhOC_3H_6OPhNH_2$). With two amine groups on the same molecule, two carboxylate groups on two polyoxometalates can form amide bonds with the same amine, tethering the two POMs together. The products of these reactions have been characterized spectroscopically.



Keggin ion with rhodium-carboxylate adduct occupying vacancy