



**Illinois Wesleyan University**  
**Digital Commons @ IWU**

---

John Wesley Powell Student Research  
Conference

2003, 14th Annual JWP Conference

---

Apr 12th, 9:00 AM - 10:00 AM

**Variation in Rates of Asexual Reproduction By *Convolutriloba*  
*Retrogemma***

Sara Rahim  
*Illinois Wesleyan University*

William Jaeckle, Faculty Advisor  
*Illinois Wesleyan University*

Follow this and additional works at: <https://digitalcommons.iwu.edu/jwprc>

---

Rahim, Sara and Jaeckle, Faculty Advisor, William, "Variation in Rates of Asexual Reproduction By *Convolutriloba Retrogemma*" (2003). *John Wesley Powell Student Research Conference*. 22.

<https://digitalcommons.iwu.edu/jwprc/2003/posters/22>

This Event is protected by copyright and/or related rights. It has been brought to you by Digital Commons @ IWU with permission from the rights-holder(s). You are free to use this material in any way that is permitted by the copyright and related rights legislation that applies to your use. For other uses you need to obtain permission from the rights-holder(s) directly, unless additional rights are indicated by a Creative Commons license in the record and/ or on the work itself. This material has been accepted for inclusion by faculty at Illinois Wesleyan University. For more information, please contact [digitalcommons@iwu.edu](mailto:digitalcommons@iwu.edu).

©Copyright is owned by the author of this document.

Poster Presentation P33

**VARIATION IN RATES OF ASEXUAL REPRODUCTION BY  
*CONVOLUTRILLOBA RETROGEMMA***

Sara Rahim and William Jaeckle\*

Department of Biology, Illinois Wesleyan University

*Convolutriloba retrogemma*, an acoel turbellarian (phylum Platyhelminthes), engages in an obligate symbiotic relationship with unicellular algae. This species reproduces asexually by budding from the posterior end of the parent individual. The rate of reproduction of 24 newly budded flatworms of various sizes was studied over a period of six weeks. Flatworms were individually placed into 6 mL of 0.2  $\mu$ m filtered seawater of 33.2 ppt salinity in each of the 6 wells of a multiwell plate. All plates were placed into an incubator set at a constant temperature of 25° C with invariable light conditions (12D:12L light cycle). Reproduction reached a peak during the third and fourth weeks when each flatworm produced an average of four clones. The average overall rate of reproduction was one clone every four to five days. However, observations of the budding process revealed a positive relationship between the size of the parent to the size of the clone and the number of clones produced, which may explain the substantial amount of variability in the rate of reproduction among individual flatworms. The results of this study suggest that the nature of the symbiotic relationship between the algae and the flatworm is mutualistic. Since budding proceeded without access to prey, the environmental conditions were adequate to promote photosynthesis by the algal symbionts resulting in sufficient energy for the flatworm to reproduce. The flatworm, in return, provides shelter and potentially nutrients to the algae.