



Illinois Wesleyan University
Digital Commons @ IWU

John Wesley Powell Student Research
Conference

2004, 15th Annual JWP Conference

Apr 17th, 1:15 PM - 2:30 PM

Designing FNR Mutant Proteins to Probe Conformational Changes

Jena Hudson

Illinois Wesleyan University

Mialy Ramarosan

Illinois Wesleyan University

Robin Johnson

Illinois Wesleyan University

Laura Moore, Faculty Advisor

Illinois Wesleyan University

Follow this and additional works at: <https://digitalcommons.iwu.edu/jwprc>

Hudson, Jena; Ramarosan, Mialy; Johnson, Robin; and Moore, Faculty Advisor, Laura, "Designing FNR Mutant Proteins to Probe Conformational Changes" (2004). *John Wesley Powell Student Research Conference*. 12.

<https://digitalcommons.iwu.edu/jwprc/2004/posters2/12>

This Event is protected by copyright and/or related rights. It has been brought to you by Digital Commons @ IWU with permission from the rights-holder(s). You are free to use this material in any way that is permitted by the copyright and related rights legislation that applies to your use. For other uses you need to obtain permission from the rights-holder(s) directly, unless additional rights are indicated by a Creative Commons license in the record and/ or on the work itself. This material has been accepted for inclusion by faculty at Illinois Wesleyan University. For more information, please contact digitalcommons@iwu.edu.

©Copyright is owned by the author of this document.

Poster Presentation P24

**DESIGNING FNR MUTANT PROTEINS TO
PROBE CONFORMATIONAL CHANGES**

Jena Hudson, Mialy Ramaroson, Robin Johnson and Laura Moore*
Department of Chemistry, Illinois Wesleyan University

The goal of our research is to understand the conformational changes that take place as a transcriptional factor protein (FNR) goes from its inactive to active form. FNR is a protein found in facultatively anaerobic bacteria. It becomes active and dimerizes when oxygen is not present. This dimerization occurs after a 4Fe-4S cluster is assembled. The dimerized FNR binds to the DNA and activates the transcription of genes necessary for the anaerobic pathways. Since traditional methods of determining protein structure, such as x-ray crystallography or NMR spectroscopy, have not been successful with FNR, we will be using tryptophan fluorescence to better define the conformational changes. The wavelength of tryptophan fluorescence emission varies upon exposure to solvent and can be used to infer the change of position of an amino acid between aerobic and anaerobic states. As a first step, tryptophan residues were inserted at different amino acid locations along the dimerization helix and near the N-terminus. The FNR mutants R48W, L34W, R140W, Q141W, Q142W, M143W, M144W, S148W, G149W, K152W, G153W and G156W were created by site-directed mutagenesis, in the pET11a plasmid. Of these mutations, Q141W, M143W, M144W, G149W and E150W retained activity similar to the wild type protein. Wild type activity, for our purposes, is defined as at least 50% the activity of the unmutated protein. The retention of wild type activity is crucial to our study because only functional proteins, which are able to dimerize properly, will give insight regarding the actual conformational changes and structure of FNR. In an additional experiment, L146W, M147W and M157W were cloned into the pACYC184 plasmid in order to investigate dimerization of the mutant at FNR concentrations similar to the concentrations likely to be used for in-vitro experiments. Of the three, L146W and M157W retained wild type activity. Currently, we are attempting to purify the mutant proteins using column chromatography.