



Illinois Wesleyan University
Digital Commons @ IWU

John Wesley Powell Student Research
Conference

2004, 15th Annual JWP Conference

Apr 17th, 1:15 PM - 2:30 PM

Functionality of FNR Proteins with Mutations in the FE-S Binding Domain

Molly Marker
Illinois Wesleyan University

Laura Moore, Faculty Advisor
Illinois Wesleyan University

Follow this and additional works at: <https://digitalcommons.iwu.edu/jwprc>

Marker, Molly and Moore, Faculty Advisor, Laura, "Functionality of FNR Proteins with Mutations in the FE-S Binding Domain" (2004). *John Wesley Powell Student Research Conference*. 14.

<https://digitalcommons.iwu.edu/jwprc/2004/posters2/14>

This Event is protected by copyright and/or related rights. It has been brought to you by Digital Commons @ IWU with permission from the rights-holder(s). You are free to use this material in any way that is permitted by the copyright and related rights legislation that applies to your use. For other uses you need to obtain permission from the rights-holder(s) directly, unless additional rights are indicated by a Creative Commons license in the record and/ or on the work itself. This material has been accepted for inclusion by faculty at Illinois Wesleyan University. For more information, please contact digitalcommons@iwu.edu.

©Copyright is owned by the author of this document.

Poster Presentation P28

**FUNCTIONALITY OF FNR PROTEINS WITH MUTATIONS IN THE FE-S
BINDING DOMAIN**

Molly Marker and Laura Moore*

Department of Biology, Illinois Wesleyan University

The FNR protein is an oxygen sensing transcription factor found in *Escherichia coli*. The binding of wildtype FNR to DNA is facilitated by the presence of a 4Fe-4S cluster, which functions to dimerize the protein. Entry of oxygen into the cell destroys the 4Fe-4S cluster and causes a conformational change in the protein, which then impedes dimerization. It is not understood how or why oxygen destroys the 4Fe-4S cluster. However, it is known that a leucine to histidine mutation at position 28 retains a stable 4Fe-4S cluster in the presence of oxygen. The properties of the amino acid residues (charge, size) surrounding the Fe-S cluster are important when considering its stability. The four cysteine amino acids found at positions 20, 23, 29 and 122 in the coding sequence of FNR ligand the four iron molecules in the cluster. We have investigated the importance of particular residues near these cysteine ligands. Our anaerobic cultures show decreased activity of FNR in alanine and aspartate mutants compared to histidine and lysine mutants at position 28. Aerobic studies are currently being conducted.