



Illinois Wesleyan University
Digital Commons @ IWU

John Wesley Powell Student Research
Conference

2005, 16th Annual JWP Conference

Apr 16th, 1:15 PM - 2:30 PM

A Taxonomic Study of the Lichen Genus *Rimelia* in the Great Smoky Mountains National Park

Erin Boente
Illinois Wesleyan University

Jonathan Dey, Faculty Advisor
Illinois Wesleyan University

Follow this and additional works at: <https://digitalcommons.iwu.edu/jwprc>

Boente, Erin and Dey, Faculty Advisor, Jonathan, "A Taxonomic Study of the Lichen Genus *Rimelia* in the Great Smoky Mountains National Park" (2005). *John Wesley Powell Student Research Conference*. 4.

<https://digitalcommons.iwu.edu/jwprc/2005/posters2/4>

This Event is protected by copyright and/or related rights. It has been brought to you by Digital Commons @ IWU with permission from the rights-holder(s). You are free to use this material in any way that is permitted by the copyright and related rights legislation that applies to your use. For other uses you need to obtain permission from the rights-holder(s) directly, unless additional rights are indicated by a Creative Commons license in the record and/ or on the work itself. This material has been accepted for inclusion by faculty at Illinois Wesleyan University. For more information, please contact digitalcommons@iwu.edu.

©Copyright is owned by the author of this document.

Poster Presentation P10

A TAXONOMIC STUDY OF THE LICHEN GENUS RIMELIA IN THE GREAT SMOKY MOUNTAINS NATIONAL PARK

Erin Boente and Jonathan Dey*

Biology Department, Illinois Wesleyan University

A lichen is a fungus that grows in a symbiotic relationship with a photosynthetic green alga or cyanobacterium. *Rimelia* species are broad-lobed, foliose lichens that have been recently segregated from *Parmotrema*. Distinctively, the former's lobes are strongly maculate (spotted) in a reticulate pattern (and are often cracked) and they produce rhizines on their lower cortex all the way to the lobe margins.

The 1997 National Park Service checklist of lichen species in the Great Smoky Mountains National Park (GSMNP) included only two *Rimelia* species (*R. reticulata* and *R. subsidiosa*). Field and laboratory work by Jonathan Dey and Illinois Wesleyan University undergraduate students (Rebecca Rincker and Holly Grey in 1998; Jana Rose and Adrienne Gagnon in 2000; Emily Richter and Sarah Mick in 2003) pointed to the presence of additional species of *Rimelia* in the GSMNP. As a result, we have undertaken an examination of all *Rimelia* specimens previously collected in the GSMNP and deposited in the lichen herbarium at Illinois Wesleyan University in order to update the species list for the Park and to produce an identification key and descriptions to all species of *Rimelia* in the GSMNP.

As a result of this study, five species of *Rimelia* are now known to occur in the GSMNP. *Rimelia cetrata*, *R. commensurata*, and *R. simulans* are newly reported in the Park to go with the previously known *R. reticulata* and *R. subsidiosa*. *Rimelia diffracta* occurs in the southern Appalachian Mountain region but has not yet been found in the Park. The additional products of this study include an identification key to *Rimelia* species and species descriptions including morphology, secondary product chemistry, local ecology, general distribution, a list of specimens examined, and GSMNP distribution map.

This study is a part of the All Taxa Biodiversity Inventory (ATBI) of the GSMNP, a large endeavor where scientists and educators are working collaboratively to determine all the organisms that can be found in the park.

The work reported here is a portion of Erin Boente's Research Honor's Project in the Biology Department at IWU. She is also working on two other genera---*Canoparmelia* and *Canomaculina*---in the GSMNP.