



Apr 16th, 9:00 AM - 10:00 AM

## Multistep Synthesis of Asymmetric Organophosphorus Analogs of Acetylcholine

Jonathan Kaplan  
*Illinois Wesleyan University*

Dan Flaughner  
*Illinois Wesleyan University*

Jeffrey Frick, Faculty Advisor  
*Illinois Wesleyan University*

Follow this and additional works at: <https://digitalcommons.iwu.edu/jwprc>

---

Kaplan, Jonathan; Flaughner, Dan; and Frick, Faculty Advisor, Jeffrey, "Multistep Synthesis of Asymmetric Organophosphorus Analogs of Acetylcholine" (2005). *John Wesley Powell Student Research Conference*. 14.

<https://digitalcommons.iwu.edu/jwprc/2005/posters/14>

This is protected by copyright and/or related rights. It has been brought to you by Digital Commons @ IWU with permission from the rights-holder(s). You are free to use this material in any way that is permitted by the copyright and related rights legislation that applies to your use. For other uses you need to obtain permission from the rights-holder(s) directly, unless additional rights are indicated by a Creative Commons license in the record and/ or on the work itself. This material has been accepted for inclusion by faculty at Illinois Wesleyan University. For more information, please contact [digitalcommons@iwu.edu](mailto:digitalcommons@iwu.edu).

©Copyright is owned by the author of this document.

Poster Presentation P23

**MULTISTEP SYNTHESIS OF ASYMMETRIC ORGANOPHOSPHORUS  
ANALOGS OF ACETYLCHOLINE**

Jonathan Kaplan, Dan Flaugh, and Jeffrey Frick\*  
Chemistry Department, Illinois Wesleyan University

Acetylcholinesterase is an enzyme which catalyzes the hydrolysis of acetylcholine, causing the relaxation of smooth, cardiac, and skeletal muscle contractions in the human body. One class of compounds that inhibit acetylcholinesterase is the organophosphates. While the mechanism of inhibition is fairly well understood, much less is known about the stereochemical aspects of inhibition. Our research involves the multi-step synthesis of organophosphorous analogs of acetylcholine which have similar size and functionality of that of acetylcholine, but that also contain two chiral centers. Once prepared, these analogs could then be used to learn more about the active site of acetylcholinesterase and also to learn about the stereochemistry involved in the phosphorylation of acetylcholinesterase. We are currently working on separating pairs of diastereomers, which will eventually lead to the production of the desired analogs.