



**Illinois Wesleyan University**  
**Digital Commons @ IWU**

---

John Wesley Powell Student Research  
Conference

2005, 16th Annual JWP Conference

---

Apr 16th, 9:00 AM - 10:00 AM

## **The Role of the Hinge Region in the Activity of the Protein FNR in E. Coli**

Katherine McCulloch  
*Illinois Wesleyan University*

Laura Moore, Faculty Advisor  
*Illinois Wesleyan University*

Follow this and additional works at: <https://digitalcommons.iwu.edu/jwprc>

---

McCulloch, Katherine and Moore, Faculty Advisor, Laura, "The Role of the Hinge Region in the Activity of the Protein FNR in E. Coli" (2005). *John Wesley Powell Student Research Conference*. 24.

<https://digitalcommons.iwu.edu/jwprc/2005/posters/24>

This Event is protected by copyright and/or related rights. It has been brought to you by Digital Commons @ IWU with permission from the rights-holder(s). You are free to use this material in any way that is permitted by the copyright and related rights legislation that applies to your use. For other uses you need to obtain permission from the rights-holder(s) directly, unless additional rights are indicated by a Creative Commons license in the record and/ or on the work itself. This material has been accepted for inclusion by faculty at Illinois Wesleyan University. For more information, please contact [digitalcommons@iwu.edu](mailto:digitalcommons@iwu.edu).

©Copyright is owned by the author of this document.

Poster Presentation P41

**THE ROLE OF THE HINGE REGION IN THE ACTIVITY OF  
THE PROTEIN FNR IN E. COLI**

Katherine McCulloch and Laura Moore\*  
Chemistry Department, Illinois Wesleyan University

FNR is an oxygen-sensing transcription factor in the facultative anaerobic bacteria *E. coli*. It is part of a family of proteins that includes cAMP receptor proteins (CRP or CAP). These proteins undergo conformational changes in response to specific effector molecules, which in the case of FNR is oxygen. In its functional form, two subunits of the FNR protein form a homodimer through a  $\alpha$ -helix, residues 140 to 160. Adjacent to this helix is a "hinge" or "switch" region at residues 161 to 171, far from the oxygen-sensing domain of the protein. In homologues to FNR, CAP and CooA, it has been shown that this region plays a role in stabilizing the conformational changes involved in the activation process. Residues 161, 169, 170, and 171 in particular are highly conserved in the CRP family. The goal of this project is to study the importance of this switch region in the activity of FNR. Site-directed mutagenesis and  $\alpha$ -galactosidase activity assays were performed to investigate the role of the hinge region in FNR, and they have shown that this region may have less involvement in the activation process than its homologues. Future work will involve reversing the charges of several residues of the hinge region, and analyzing how such a change affects the activity of FNR..