



Apr 17th, 9:00 AM - 10:00 AM

## Isolation and Characterization of Gibberellins from the Moss *C. Purpureus*

Alison M. Redden  
*Illinois Wesleyan University*

Rebecca Roesner, Faculty Advisor  
*Illinois Wesleyan University*

Loni Walker, Faculty Advisor  
*Illinois Wesleyan University*

Follow this and additional works at: <https://digitalcommons.iwu.edu/jwprc>

---

Redden, Alison M.; Roesner, Faculty Advisor, Rebecca; and Walker, Faculty Advisor, Loni, "Isolation and Characterization of Gibberellins from the Moss *C. Purpureus*" (2004). *John Wesley Powell Student Research Conference*. 24.  
<https://digitalcommons.iwu.edu/jwprc/2004/posters/24>

This Event is protected by copyright and/or related rights. It has been brought to you by Digital Commons @ IWU with permission from the rights-holder(s). You are free to use this material in any way that is permitted by the copyright and related rights legislation that applies to your use. For other uses you need to obtain permission from the rights-holder(s) directly, unless additional rights are indicated by a Creative Commons license in the record and/ or on the work itself. This material has been accepted for inclusion by faculty at Illinois Wesleyan University. For more information, please contact [digitalcommons@iwu.edu](mailto:digitalcommons@iwu.edu).

©Copyright is owned by the author of this document.


Poster Presentation P43

ISOLATION AND CHARACTERIZATION OF GIBBERELLINS  
FROM THE MOSS *C. PURPUREUS*

Alison M. Redden and Rebecca Roesner\* and Loni Walker \*  
Department of Chemistry and Biology, Illinois Wesleyan University

The plant hormone Gibberellic Acid (GA) is known to mediate a number of developmental processes in flowering plants, including fruit growth, germination, and internode elongation. Although the information concerning GAs in flowering plants is well documented, very few studies have addressed the roles of GAs in non-flowering bryophytes, specifically the mosses.

Recent evidence suggests that GA may also play an important role in the growth and development of mosses, specifically *Ceratodon purpureus*. Preliminary studies utilizing a GA biosynthesis inhibitor caused a dramatic decrease in *C. purpureus* growth, while the application of exogenous GA restored normal growth. These observations lead to the conclusion that Gibberellic Acid is vital to normal growth and function in the mosses. Results from further growth studies, as well as current attempts to isolate and characterize gibberellins from plant tissue are reported.



GA<sub>4</sub> – Commercially available C<sub>19</sub> GA