



Illinois Wesleyan University
Digital Commons @ IWU

John Wesley Powell Student Research
Conference

2005, 16th Annual JWP Conference

Apr 16th, 9:00 AM - 10:00 AM

Electrophysiological Study of the Anterior Cingulate

Mark N. Rubin
Illinois Wesleyan University

Joseph Williams, Faculty Advisor
Illinois Wesleyan University

Follow this and additional works at: <https://digitalcommons.iwu.edu/jwprc>

Rubin, Mark N. and Williams, Faculty Advisor, Joseph, "Electrophysiological Study of the Anterior Cingulate" (2005). *John Wesley Powell Student Research Conference*. 28.
<https://digitalcommons.iwu.edu/jwprc/2005/posters/28>

This Event is protected by copyright and/or related rights. It has been brought to you by Digital Commons @ IWU with permission from the rights-holder(s). You are free to use this material in any way that is permitted by the copyright and related rights legislation that applies to your use. For other uses you need to obtain permission from the rights-holder(s) directly, unless additional rights are indicated by a Creative Commons license in the record and/ or on the work itself. This material has been accepted for inclusion by faculty at Illinois Wesleyan University. For more information, please contact digitalcommons@iwu.edu.

©Copyright is owned by the author of this document.

Poster Presentation P47

ELECTROPHYSIOLOGICAL STUDY OF THE ANTERIOR CINGULATE

Mark N. Rubin and Joseph Williams*
Psychology Department, Illinois Wesleyan University

Theta EEG activity has been well-characterized in the hippocampus, a medial temporal lobe structure essential for learning and memory. For instance, it is known that the medial septal area (MSA) drives hippocampal theta rhythmicity. However, comparatively little is known about the neuroanatomical substrates behind theta activity in other brain areas, such as the anterior cingulate region of the prefrontal cortex. One aim of this study is to address this question by determining whether or not the MSA drives theta rhythmicity in the anterior cingulate as it does in the hippocampus. To meet this aim, EEG recording electrodes will be surgically implanted into the right anterior cingulate (AC) region and a guide cannula will be placed in the MSA of male Long-Evans rats. Once the rats have fully recovered from surgery, the investigators will monitor the electrophysiological behavior of the AC. Once theta rhythmicity is established and recorded, the investigators will induce a temporary lesion in the MSA while simultaneously recording AC theta to determine whether a lesion to the medial septal area abolishes the AC theta rhythm. Experimental results will be discussed at the conference.