



Apr 8th, 10:00 AM - 11:00 AM

Characterization of a TIR-like Gene in the Moss *Physcomitrella Patens*

Rachael Liesman
Illinois Wesleyan University

Loni Walker, Faculty Advisor
Illinois Wesleyan University

Follow this and additional works at: <https://digitalcommons.iwu.edu/jwprc>

Liesman, Rachael and Walker, Faculty Advisor, Loni, "Characterization of a TIR-like Gene in the Moss *Physcomitrella Patens*" (2006). *John Wesley Powell Student Research Conference*. 3.

<https://digitalcommons.iwu.edu/jwprc/2006/oralpres3/3>

This is protected by copyright and/or related rights. It has been brought to you by Digital Commons @ IWU with permission from the rights-holder(s). You are free to use this material in any way that is permitted by the copyright and related rights legislation that applies to your use. For other uses you need to obtain permission from the rights-holder(s) directly, unless additional rights are indicated by a Creative Commons license in the record and/ or on the work itself. This material has been accepted for inclusion by faculty at Illinois Wesleyan University. For more information, please contact digitalcommons@iwu.edu.

©Copyright is owned by the author of this document.

Oral Presentation O3.3

**CHARACTERIZATION OF A TIR-LIKE GENE IN THE MOSS
PHYSCOMITRELLA PATENS**

Rachael Liesman and Loni Walker*
Biology Department, Illinois Wesleyan University

Auxin is a plant hormone mediating a wide variety of plant developmental and growth processes. One way that auxin exerts these effects is by regulating gene expression.

Aux/IAA proteins are transcription repressors that interact with specific transcription factors known as ARFs. TIR and TIR-like genes encode f-box proteins that promote the ubiquitin-mediated degradation of Aux/IAA proteins and thus are also important regulators of plant development. In the presence of increased auxin concentrations, auxin binds to TIR1, promoting its association with the SCF complex. TIR1 then targets Aux/IAA proteins for ubiquitination by the SCF complex. This results in the degradation of Aux/IAA proteins and the release of ARFs. ARFs are then free to promote transcription of downstream genes important in plant growth and development.

I am interested in understanding TIR function in mosses. One TIR-like gene encoded in the moss *Physcomitrella patens* genome is 18f14. Using cloning, PCR, and sequencing techniques, I have characterized the gene structure and isolated 3 introns within 18f14.