



Illinois Wesleyan University
Digital Commons @ IWU

John Wesley Powell Student Research
Conference

2006, 17th Annual JWP Conference

Apr 8th, 9:00 AM - 10:00 AM

Oxygen Sensing in the FNR Transcription Factor: Creation and Characterization of Variants with Aerobically Stable [4FE-4S] Clusters

Robin Johnson
Illinois Wesleyan University

Maura Bates
Illinois Wesleyan University

Laura Moore, Faculty Advisor
Illinois Wesleyan University

Follow this and additional works at: <https://digitalcommons.iwu.edu/jwprc>

Johnson, Robin; Bates, Maura; and Moore, Faculty Advisor, Laura, "Oxygen Sensing in the FNR Transcription Factor: Creation and Characterization of Variants with Aerobically Stable [4FE-4S] Clusters" (2006). *John Wesley Powell Student Research Conference*. 40. <https://digitalcommons.iwu.edu/jwprc/2006/posters/40>

This Event is protected by copyright and/or related rights. It has been brought to you by Digital Commons @ IWU with permission from the rights-holder(s). You are free to use this material in any way that is permitted by the copyright and related rights legislation that applies to your use. For other uses you need to obtain permission from the rights-holder(s) directly, unless additional rights are indicated by a Creative Commons license in the record and/ or on the work itself. This material has been accepted for inclusion by faculty at Illinois Wesleyan University. For more information, please contact digitalcommons@iwu.edu.

©Copyright is owned by the author of this document.

Poster Presentation P7

**OXYGEN SENSING IN THE FNR TRANSCRIPTION FACTOR: CREATION
AND CHARACTERIZATION OF VARIANTS WITH AEROBICALLY STABLE
[4Fe-4S] CLUSTERS**

Robin Johnson, and Maura Bates and Laura Moore*
Chemistry Department, Illinois Wesleyan University

Activity of the FNR protein, a transcription factor in the facultative anaerobe *Escherichia coli*, is contingent upon the integrity of a [4Fe-4S] cluster incorporated by the protein's effector domain. The cluster is ligated by cysteine residues at positions 20, 23, 29, and 122. In wild-type FNR, this [4Fe-4S] cluster degrades upon exposure to molecular oxygen, and the resulting conformational changes cause the homodimer to separate into monomers, deactivating the protein. A previous study has identified an FNR variant, Leu28His, which has an oxygen-stable [4Fe-4S] cluster. This mutant protein retains activity under aerobic conditions. Using beta-galactosidase assays to quantify in vivo FNR activity, our lab has explored the oxygen sensitivity of several other FNR variants with amino acid substitutions adjacent to ligating cysteines. While many of these proteins exhibit oxygen sensitivity similar to that of wild-type FNR, the mutants Leu28Lys, Leu28Asn, Leu28Gln, Leu28Ser, and Leu28Tyr all retain some activity under aerobic conditions. Western blot analysis suggests that this increase in activity as compared to wild-type is not due to an increase in protein concentration. These mutant proteins have been isolated and characterized by absorption spectroscopy. The spectra indicate that the [4Fe-4S] clusters of these FNR variants do exhibit some degree of oxygen stability. As only substitutions of amino acids with amine, alcohol, or amide side chains have so far resulted in aerobically active FNR, oxygen stability of the [Fe-4S] cluster may result from donation of a hydrogen bond at position 28.