



Apr 8th, 9:00 AM - 10:00 AM

Organochlorine Pesticide Contamination in Resident North American Passerines

Amy Cadwallader
Illinois Wesleyan University

Kevin Latman,
Illinois Wesleyan University

Given Harper, Faculty Advisor
Illinois Wesleyan University

Jeffrey Frick, Faculty Advisor
Illinois Wesleyan University

Follow this and additional works at: <https://digitalcommons.iwu.edu/jwprc>

Cadwallader, Amy; Latman,, Kevin; Harper, Faculty Advisor, Given; and Frick, Faculty Advisor, Jeffrey, "Organochlorine Pesticide Contamination in Resident North American Passerines" (2006). *John Wesley Powell Student Research Conference*. 6.
<https://digitalcommons.iwu.edu/jwprc/2006/posters/6>

This is protected by copyright and/or related rights. It has been brought to you by Digital Commons @ IWU with permission from the rights-holder(s). You are free to use this material in any way that is permitted by the copyright and related rights legislation that applies to your use. For other uses you need to obtain permission from the rights-holder(s) directly, unless additional rights are indicated by a Creative Commons license in the record and/ or on the work itself. This material has been accepted for inclusion by faculty at Illinois Wesleyan University. For more information, please contact digitalcommons@iwu.edu.

©Copyright is owned by the author of this document.

Poster Presentation P15

**ORGANOCHLORINE PESTICIDE CONTAMINATION IN RESIDENT
NORTH AMERICAN PASSERINES**

Amy Cadwallader and Kevin Latman and Given Harper and Jeffrey Frick*
Biology and Chemistry Departments, Illinois Wesleyan University

Organochlorine (OC) pesticide and metabolite levels were determined in resident North American passerines and woodpeckers collected in McLean County, Illinois. Sixty-nine of 77 individual birds contained OC compounds above detection limits, including all eighteen species examined. Total contamination levels in individual birds ranged from 7.47 to 2,274.23 ng/g. The most prevalent OC compound was p,p'-DDE, which was present in 45 birds; dieldrin was found in 30 birds, heptachlor and heptachlor epoxide were each found in 27 birds, and p,p'-DDT was found in 25 birds. There was no significant difference in total OC levels between males and females, between HY and AHY birds, and between birds that bred and wintered in more northern latitudes. There was a significant difference in total OC levels between diet categories. Omnivores had significantly higher total OC levels than granivores. House sparrows had significantly lower total OC levels than both white-throated sparrows and American robins.