



Apr 12th, 9:00 AM - 10:00 AM

Caudal Fin Skeletal Development: A Comparison Between *Moenkhausia Sanctaefilomenae* and *Danio Rerio*

Mira Coleman
Illinois Wesleyan University

Brian Walter, Faculty Advisor
Illinois Wesleyan University

Follow this and additional works at: <https://digitalcommons.iwu.edu/jwprc>

Coleman, Mira and Walter, Faculty Advisor, Brian, "Caudal Fin Skeletal Development: A Comparison Between *Moenkhausia Sanctaefilomenae* and *Danio Rerio*" (2008). *John Wesley Powell Student Research Conference*. 5.

<https://digitalcommons.iwu.edu/jwprc/2008/posters/5>

This Event is protected by copyright and/or related rights. It has been brought to you by Digital Commons @ IWU with permission from the rights-holder(s). You are free to use this material in any way that is permitted by the copyright and related rights legislation that applies to your use. For other uses you need to obtain permission from the rights-holder(s) directly, unless additional rights are indicated by a Creative Commons license in the record and/ or on the work itself. This material has been accepted for inclusion by faculty at Illinois Wesleyan University. For more information, please contact digitalcommons@iwu.edu.

©Copyright is owned by the author of this document.

Poster Presentation P9

**CAUDAL FIN SKELETAL DEVELOPMENT: A COMPARISON BETWEEN
MOENKHAUSIA SANCTAEFILOMENAE AND *DANIO RERIO***

Mira Coleman and Brian Walter*
Biology Department, Illinois Wesleyan University

Our research is focused on elucidating the developmental mechanisms that underlie morphological diversity among closely related organisms. In this study, embryological and histological techniques were used to analyze developmental variation in the caudal fin between two species of fishes, *Moenkhausia sanctaefilomenae* and *Danio rerio*. To visualize the patterning of cartilage and bone growth in the caudal fin, whole mount Alizarin red and Alcian blue staining was used to bind calcium and cartilage extracellular matrix, respectively. Thus, the dynamic processes of cartilage growth and endochondral ossification could be visualized at various larval developmental time-points. When comparing the *M. sanctaefilomenae* and *D. rerio*, time-points that are correlated with significant changes in the developing caudal fin could be established and used for frames of reference. In general, *M. sanctaefilomenae* displayed more rapid skeletal development in the caudal fin than did *D. rerio*. The next step of our research will include the analysis of genes known to be responsible for skeletal development in order to account for the differences observed between these two species.