



Illinois Wesleyan University
Digital Commons @ IWU

John Wesley Powell Student Research
Conference

2003, 14th Annual JWP Conference

Apr 12th, 1:15 PM - 2:15 PM

Development of Amine Protecting Group Strategies Compatible with the Hexamolybdate Ion

Elizabeth K. Myers
Illinois Wesleyan University

Rebecca Roesner, Faculty Advisor
Illinois Wesleyan University

Follow this and additional works at: <https://digitalcommons.iwu.edu/jwprc>

Myers, Elizabeth K. and Roesner, Faculty Advisor, Rebecca, "Development of Amine Protecting Group Strategies Compatible with the Hexamolybdate Ion" (2003). *John Wesley Powell Student Research Conference*. 12.
<https://digitalcommons.iwu.edu/jwprc/2003/posters2/12>

This Event is protected by copyright and/or related rights. It has been brought to you by Digital Commons @ IWU with permission from the rights-holder(s). You are free to use this material in any way that is permitted by the copyright and related rights legislation that applies to your use. For other uses you need to obtain permission from the rights-holder(s) directly, unless additional rights are indicated by a Creative Commons license in the record and/ or on the work itself. This material has been accepted for inclusion by faculty at Illinois Wesleyan University. For more information, please contact digitalcommons@iwu.edu.

©Copyright is owned by the author of this document.

Poster Presentation P28

DEVELOPMENT OF AMINE PROTECTING GROUP STRATEGIES
COMPATIBLE WITH THE HEXAMOLYBDATE ION

Elizabeth K. Myers and Rebecca Roesner*

Department of Chemistry, Illinois Wesleyan University

Polyoxometalates are large, symmetrical, anionic metal-oxygen clusters of the groups V and VI transition metals in their highest oxidation states. They are known for their many interesting and useful properties: catalytic activity, reversible oxidation, applications in supramolecular chemistry, and anti-viral and anti-tumoral behaviors. Many of these properties could be better utilized through the introduction of organic and bio-organic substituents as linkers, handles, or tethers to the surface of the polyoxometalate. One widely studied polyoxometalate is the hexamolybdate ion $[\text{Mo}_6\text{O}_{19}]^{2-}$. One possible tool to better control the products of reactions between aromatic diamines and $[\text{n-Bu}_4\text{N}]_2[\text{Mo}_6\text{O}_{19}]$ is to use protecting groups.

A method was developed to protect only one end of a difunctional amine with the BOC protecting group. The BOC protecting group was successfully used to protect 1,4-butylene-bis(phenoxy-4-amine), and mono-BOC-protected 1,4-butylene-bis(phenoxy-4-amine) was isolated. The mono-BOC-protected 1,4-butylene-bis(phenoxy-4-amine) was then successfully reacted with $[\text{n-Bu}_4\text{N}]_2[\text{Mo}_6\text{O}_{19}]$ (Figure 1).




Figure 1: The reaction of mono-BOC-protected 1,4-butylene-bis(phenoxy-4-amine) with $[\text{n-Bu}_4\text{N}]_2[\text{Mo}_6\text{O}_{19}]$