



Illinois Wesleyan University
Digital Commons @ IWU

John Wesley Powell Student Research
Conference

2003, 14th Annual JWP Conference

Apr 12th, 1:15 PM - 2:15 PM

The Synthesis and Isolation of Mono-BOC-Protected 1,4-Phenylene Diamine and Further Reaction with Hexamolybdate

Matthew S. Ralph
Illinois Wesleyan University

Rebecca Roesner, Faculty Advisor
Illinois Wesleyan University

Follow this and additional works at: <https://digitalcommons.iwu.edu/jwprc>

Ralph, Matthew S. and Roesner, Faculty Advisor, Rebecca, "The Synthesis and Isolation of Mono-BOC-Protected 1,4-Phenylene Diamine and Further Reaction with Hexamolybdate" (2003). *John Wesley Powell Student Research Conference*. 9.
<https://digitalcommons.iwu.edu/jwprc/2003/posters2/9>

This Event is protected by copyright and/or related rights. It has been brought to you by Digital Commons @ IWU with permission from the rights-holder(s). You are free to use this material in any way that is permitted by the copyright and related rights legislation that applies to your use. For other uses you need to obtain permission from the rights-holder(s) directly, unless additional rights are indicated by a Creative Commons license in the record and/ or on the work itself. This material has been accepted for inclusion by faculty at Illinois Wesleyan University. For more information, please contact digitalcommons@iwu.edu.

©Copyright is owned by the author of this document.

Poster Presentation P34

**THE SYNTHESIS AND ISOLATION OF MONO-BOC-PROTECTED
1,4-PHENYLENE DIAMINE AND FURTHER REACTION WITH
HEXAMOLYBDATE**

Matthew S. Ralph and Rebecca A. Roesner*
Department of Chemistry, Illinois Wesleyan University

When reacting unprotected diamines with polyoxymetalates, a mixture of di-, mono-, and unsubstituted products is obtained. A simple aromatic diamine, 1,4-phenylene diamine, was used to investigate a method through which a mono-BOC-protected diamine could be synthesized and isolated for further reaction with the hexamolybdate ion to obtain a mono-substituted product. The 1,4-phenylene diamine was purified by sublimation and the purified diamine was then reacted in a 1:1 mole ratio with di-*tert*-butyl dicarbonate to mono-protect the diamine. According to GC-Mass Spectrometry, the acetonitrile-soluble products of this reaction were 97.7% mono-protected and 2.0% di-protected. The mono-protected product was then purified via column chromatography with final percent yields of 89.1% and 81.5% in two trials. Purity was verified by GC-Mass Spectrometry and H-NMR. The mono-protected 1,4-phenylene diamine was reacted with the hexamolybdate ion to yield a mono-substituted 1,4-phenylene diamine.