



Illinois Wesleyan University
Digital Commons @ IWU

John Wesley Powell Student Research
Conference

1997, 8th Annual JWP Conference

Apr 12th, 10:45 AM - 11:00 AM

Adapting Daubechies' Wavelet Iteration Scheme to Biorthogonal Scaling Functions

Nathan Mueggenburg
Illinois Wesleyan University

Tian-Xiao He, Faculty Advisor
Illinois Wesleyan University

Follow this and additional works at: <https://digitalcommons.iwu.edu/jwprc>

Mueggenburg, Nathan and He, Faculty Advisor, Tian-Xiao, "Adapting Daubechies' Wavelet Iteration Scheme to Biorthogonal Scaling Functions" (1997). *John Wesley Powell Student Research Conference*. 4.

<https://digitalcommons.iwu.edu/jwprc/1997/oralpres2/4>

This Event is protected by copyright and/or related rights. It has been brought to you by Digital Commons @ IWU with permission from the rights-holder(s). You are free to use this material in any way that is permitted by the copyright and related rights legislation that applies to your use. For other uses you need to obtain permission from the rights-holder(s) directly, unless additional rights are indicated by a Creative Commons license in the record and/ or on the work itself. This material has been accepted for inclusion by faculty at Illinois Wesleyan University. For more information, please contact digitalcommons@iwu.edu.

©Copyright is owned by the author of this document.

Oral Presentation 2.2

ADAPTING DAUBECHIES' WAVELET ITERATION SCHEME TO BIORTHOGONAL SCALING FUNCTIONS

Nathan Mueggenburg and Tian-Xiao He*, Department of Mathematics, IWU

In 1988 Daubechies developed an iteration scheme to produce a D4 wavelet function which has since become very popular. This iteration scheme is based upon four conditions stated by Daubechies. One of these conditions required that the scaling function be orthogonal. We propose a new iteration scheme, similar in nature to Daubechies', but with a relaxed orthogonality condition. Our iteration scheme does not require that the scaling function be orthogonal, but only that it be biorthogonal. With this orthogonality condition relaxed, we can construct a dual of an unorthogonal function by using macroelement analysis, and thus create a larger set of possible initial functions for the iteration scheme. This new iteration scheme is then able to produce biorthogonal wavelet functions, a much broader group than that produced by Daubechies' iteration scheme. As an example we use the second order B-spline, commonly known as the hat function, as an initial function of our iteration scheme.