



Illinois Wesleyan University
Digital Commons @ IWU

John Wesley Powell Student Research
Conference

1997, 8th Annual JWP Conference

Apr 12th, 11:30 AM - 11:45 AM

Steiner Trees over Generalized Checkerboards

Meta Voelker
Illinois Wesleyan University

Robin Sanders, Faculty Advisor
Illinois Wesleyan University

Follow this and additional works at: <https://digitalcommons.iwu.edu/jwprc>

Voelker, Meta and Sanders, Faculty Advisor, Robin, "Steiner Trees over Generalized Checkerboards" (1997). *John Wesley Powell Student Research Conference*. 2.
<https://digitalcommons.iwu.edu/jwprc/1997/oralpres2/2>

This Event is protected by copyright and/or related rights. It has been brought to you by Digital Commons @ IWU with permission from the rights-holder(s). You are free to use this material in any way that is permitted by the copyright and related rights legislation that applies to your use. For other uses you need to obtain permission from the rights-holder(s) directly, unless additional rights are indicated by a Creative Commons license in the record and/ or on the work itself. This material has been accepted for inclusion by faculty at Illinois Wesleyan University. For more information, please contact digitalcommons@iwu.edu.

©Copyright is owned by the author of this document.

Oral Presentation 2.5

STEINER TREES OVER GENERALIZED CHECKERBOARDS

Meta Voelker and Robin Sanders*, Department of Mathematics, IWU

Given n randomly placed points in the plane, how can these points be connected by a network of minimum total length? Using graph theory, the answer lies in the formation of a minimum spanning tree. If additional points can be added to the tree, then the minimum spanning tree can often be shortened even further, resulting in what is known as a Steiner minimal tree. Unfortunately, no simple, general algorithm exists for finding Steiner minimal trees.

In examining a specific class of graphs, Chung, Gardner and Graham investigated building Steiner trees over "checkerboards" built of unit squares ("Steiner Trees on a Checkerboard", Mathematics Magazine, V. 62, # 2, 1989). In generalizing their ideas to rhombuses, we have discovered that for certain checkerboards with "small enough" angle, the length of their corresponding networks can be shortened by considering the checkerboards as built from isosceles triangles rather than from rhombuses.