



Apr 12th, 2:35 PM - 3:35 PM

The Role of Medial Septal Gabaergic Neurons in Prefrontal Cortex Theta Modulation

Ryan Peterson
Illinois Wesleyan University

Joseph Williams, Faculty Advisor
Illinois Wesleyan University

Follow this and additional works at: <https://digitalcommons.iwu.edu/jwprc>

Peterson, Ryan and Williams, Faculty Advisor, Joseph, "The Role of Medial Septal Gabaergic Neurons in Prefrontal Cortex Theta Modulation" (2008). *John Wesley Powell Student Research Conference*. 20.

<https://digitalcommons.iwu.edu/jwprc/2008/posters2/20>

This Event is protected by copyright and/or related rights. It has been brought to you by Digital Commons @ IWU with permission from the rights-holder(s). You are free to use this material in any way that is permitted by the copyright and related rights legislation that applies to your use. For other uses you need to obtain permission from the rights-holder(s) directly, unless additional rights are indicated by a Creative Commons license in the record and/ or on the work itself. This material has been accepted for inclusion by faculty at Illinois Wesleyan University. For more information, please contact digitalcommons@iwu.edu.

©Copyright is owned by the author of this document.

Poster Presentation P38

**THE ROLE OF MEDIAL SEPTAL GABAERGIC NEURONS IN PREFRONTAL
CORTEX THETA MODULATION**

Ryan Peterson and Joseph Williams*
Psychology Department, Illinois Wesleyan University

Theta activity recorded in various structures of the brain including the hippocampus, amygdala and prefrontal cortex (PFC) has been associated with attention, memory, decision-making and emotions. While several researchers have determined that hippocampal theta is gated by medial septal area (MSA) neuronal firing, the same has not been equally demonstrated in the PFC. Previous research in this lab has shown MSA cholinergic neurons modulate PFC theta. However it is possible that other neurotransmitter systems, such as the MSA GABAergic system also help regulate PFC theta. Given that hippocampal theta is gated by both acetylcholine and GABA, the GABAergic connections from the MSA to the PFC must be determined. Eight male Long Evans rats were used in the present experiments. A guide cannula was surgically placed in the MSA and a recording electrode was placed in the PFC. Several days later the rats were anesthetized with an injection of ketamine/xylazine and then infused with various doses of ethanol and the GABAergic agonist muscimol. The researchers predicted that both ethanol and muscimol in the MSA will reduce or eliminate theta activity in the PFC, similar to their effects on hippocampal theta.